

Some constructions of asymptotically optimal cyclic subspace codes

Chiara Castello

Università degli Studi della Campania "L. Vanvitelli"

A joint work with Paolo Santonastaso (Politecnico di Bari)



Irsee 2025

September 4, 2025

Subspace Codes

R. Koetter, F.R. Kschischang. Coding for errors and erasures in random network coding, *IEEE Transaction on Information Theory*, **2008**.

Constant dimension Subspace Codes

$\mathcal{G}_q(n, k)$: the set of k -dimensional $\mathbb{F}_q$ -subspaces of $\mathbb{F}_{q^n}$

Constant dimension Subspace Codes

$\mathcal{G}_q(n, k)$: the set of k -dimensional $\mathbb{F}_q$ -subspaces of $\mathbb{F}_{q^n}$

$\mathcal{C} \subseteq \mathcal{G}_q(n, k)$, $(\mathcal{C}, d)$ Constant dimension subspace code

$S, T \in \mathcal{G}_q(n, k)$ $d(S, T) = 2k - 2 \dim_{\mathbb{F}_q}(S \cap T)$ ***Subspace distance***

Constant dimension Subspace Codes

$\mathcal{G}_q(n, k)$: the set of k -dimensional $\mathbb{F}_q$ -subspaces of $\mathbb{F}_{q^n}$

$\mathcal{C} \subseteq \mathcal{G}_q(n, k)$, $(\mathcal{C}, d)$ Constant dimension subspace code

$S, T \in \mathcal{G}_q(n, k) \quad d(S, T) = 2k - 2 \dim_{\mathbb{F}_q}(S \cap T) \quad \textbf{Subspace distance}$

$d(\mathcal{C}) := \min\{d(S, T) : S, T \in \mathcal{C}, S \neq T\}$

Constant dimension Subspace Codes

$\mathcal{G}_q(n, k)$: the set of k -dimensional $\mathbb{F}_q$ -subspaces of $\mathbb{F}_{q^n}$

$\mathcal{C} \subseteq \mathcal{G}_q(n, k)$, $(\mathcal{C}, d)$ Constant dimension subspace code

$S, T \in \mathcal{G}_q(n, k)$ $d(S, T) = 2k - 2 \dim_{\mathbb{F}_q}(S \cap T)$ **Subspace distance**

$d(\mathcal{C}) := \min\{d(S, T) : S, T \in \mathcal{C}, S \neq T\}$

$$2 \leq d(\mathcal{C}) \leq 2k$$

Constant dimension Subspace Codes

$\mathcal{G}_q(n, k)$: the set of k -dimensional $\mathbb{F}_q$ -subspaces of $\mathbb{F}_{q^n}$

$\mathcal{C} \subseteq \mathcal{G}_q(n, k)$, $(\mathcal{C}, d)$ Constant dimension subspace code

$S, T \in \mathcal{G}_q(n, k) \quad d(S, T) = 2k - 2 \dim_{\mathbb{F}_q}(S \cap T) \quad \textbf{Subspace distance}$

$d(\mathcal{C}) := \min\{d(S, T) : S, T \in \mathcal{C}, S \neq T\}$

$$2 \leq d(\mathcal{C}) \leq 2k$$

parameters of $\mathcal{C} \quad [n, d(\mathcal{C}), |\mathcal{C}|, k]_q$

Cyclic subspace codes

$\mathcal{C} \subseteq \mathcal{G}_q(n, k)$ constant dimension subspace code in $\mathbb{F}_{q^n}$

- **T. Etzion and A. Vardy.** Error-correcting codes in projective space, *IEEE Transaction on Information Theory*, **2011**.

Cyclic subspace codes

$\mathcal{C} \subseteq \mathcal{G}_q(n, k)$ constant dimension subspace code in $\mathbb{F}_{q^n}$

$\mathcal{C}$ is **cyclic** if

$$\mathcal{C} = \cup_V \text{Orb}(V) = \cup_V \{\alpha V : \alpha \in \mathbb{F}_{q^n}^*\}$$

- **T. Etzion and A. Vardy.** Error-correcting codes in projective space, *IEEE Transaction on Information Theory*, **2011**.

Cyclic subspace codes

$$\mathcal{C} = \cup_V \text{Orb}(V) = \cup_V \{\alpha V : \alpha \in \mathbb{F}_{q^n}^*\}$$

cyclic subspace code

* **Minimum distance**

$$2 \leq d(\mathcal{C}) \leq 2k$$

* $d = 2k$

Cyclic subspace codes

$$\mathcal{C} = \cup_V \text{Orb}(V) = \cup_V \{ \alpha V : \alpha \in \mathbb{F}_{q^n}^* \}$$

cyclic subspace code

* Minimum distance

$$2 \leq d(\mathcal{C}) \leq 2k$$

* $d = 2k \Rightarrow k \mid n$

$$\mathcal{C} = \text{Orb}(\mathbb{F}_{q^k})$$

$$|\mathcal{C}| = \frac{q^n - 1}{q^k - 1}$$

F. Manganiello, E. Gorla & J. Rosenthal. Spread codes and spread decoding in network coding. In *2008 IEEE International Symposium on Information Theory* (pp. 881-885). IEEE, **2008**.

Cyclic subspace codes

$$\mathcal{C} = \cup_V \text{Orb}(V) = \cup_V \{ \alpha V : \alpha \in \mathbb{F}_{q^n}^* \}$$

cyclic subspace code

* Minimum distance

$$2 \leq d(\mathcal{C}) \leq 2k$$

$$* d = 2k \Rightarrow k \mid n$$

$$\mathcal{C} = \text{Orb}(\mathbb{F}_{q^k})$$

$$|\mathcal{C}| = \frac{q^n - 1}{q^k - 1}$$

$$* d = 2k - 2$$

$$\mathcal{C} = \text{Orb}(S) \Rightarrow |\mathcal{C}| = \frac{q^n - 1}{q - 1}$$

Cyclic subspace codes

$$\mathcal{C} = \cup_V \text{Orb}(V) = \cup_V \{\alpha V : \alpha \in \mathbb{F}_{q^n}^*\}$$

cyclic subspace code

*** Cardinality**

$\mathcal{A}_q(n, d, k) := \max \# \text{ of codewords a subspace code } \mathcal{C} \subseteq \mathcal{G}_q(n, k) \text{ may have}$

Bounds on the size of constant-dimension subspace codes

$\mathcal{A}_q(n, d, k) := \max \# \text{ of codewords a subspace code } \mathcal{C} \subseteq \mathcal{G}_q(n, k) \text{ may have}$

Sphere-packing bound

$$\mathcal{A}_q(n, d, k) \leq \left\lfloor \frac{\begin{bmatrix} n \\ k \end{bmatrix}_q}{\sum_{i=0}^{\lfloor (d/2-1)/2 \rfloor} \begin{bmatrix} k \\ i \end{bmatrix}_q \cdot \begin{bmatrix} n-k \\ i \end{bmatrix}_q \cdot q^{i^2}} \right\rfloor$$

Singleton bound

$$\mathcal{A}_q(n, d, k) \leq \begin{bmatrix} n - d/2 + 1 \\ \max\{k, n - k\} \end{bmatrix}_q$$

R. Koetter, F.R. Kschischang. Coding for errors and erasures in random network coding, *IEEE Transaction on Information Theory*, 2008.

Anticode bound

$$\mathcal{A}_q(n, d, k) \leq \frac{\begin{bmatrix} n \\ k \end{bmatrix}_q}{\begin{bmatrix} \max\{n - k, k\} + d/2 - 1 \\ d/2 - 1 \end{bmatrix}_q}$$

R. Ahlswede, H. K. Aydinian, L. H. Khachatrian. On perfect codes and related concepts. *Designs, Codes and Cryptography*, 22(3), 221-237, 2001.

H. Wang, C. Xing, R. Safavi-Naini. Linear authentication codes: bounds and constructions. *IEEE Transactions on Information Theory*, 49(4), 866-872, 2003.

Bounds on the size of constant-dimension subspace codes

$\mathcal{A}_q(n, d, k) := \max \# \text{ of codewords a subspace code } \mathcal{C} \subseteq \mathcal{G}_q(n, k) \text{ may have}$

Sphere-packing bound

$$\mathcal{A}_q(n, d, k) \leq \left\lfloor \frac{\begin{bmatrix} n \\ k \end{bmatrix}_q}{\sum_{i=0}^{\lfloor (d/2-1)/2 \rfloor} \begin{bmatrix} k \\ i \end{bmatrix}_q \cdot \begin{bmatrix} n-k \\ i \end{bmatrix}_q \cdot q^{i^2}} \right\rfloor$$

Singleton bound

R. Koetter, F.R. Kschischang. Coding for errors and erasures. *IEEE Transactions on Information Theory*, 2008.

S. Kurz. Constructions and bounds for subspace codes. *arXiv preprint arXiv:2112.11766*, 2021.

Anticode

$$\mathcal{A}_q(n, d, k) \leq \frac{\begin{bmatrix} k \\ k \end{bmatrix}_q}{\begin{bmatrix} \max\{n-k, k\} + d/2 - 1 \\ d/2 - 1 \end{bmatrix}_q}$$

R. Ahlswede, H. K. Aydinian, L. H. Khachatrian. On perfect codes and related concepts. *Designs, Codes and Cryptography*, 22(3), 221-237, 2001.

H. Wang, C. Xing, R. Safavi-Naini. Linear authentication codes: bounds and constructions. *IEEE Transactions on Information Theory*, 49(4), 866-872, 2003.

Cyclic subspace codes

$$\mathcal{C} = \cup_V \text{Orb}(V) = \cup_V \{ \alpha V : \alpha \in \mathbb{F}_{q^n}^* \}$$

cyclic subspace code

* Cardinality

$\mathcal{A}_q(n, d, k) := \max \# \text{ of codewords a subspace code } \mathcal{C} \subseteq \mathcal{G}_q(n, k) \text{ may have}$

Johnson type bound II

$$\mathcal{A}_q(n, d, k) \leq \left\lfloor \frac{q^n - 1}{q^k - 1} \left\lfloor \frac{q^{n-1} - 1}{q^{k-1} - 1} \left\lfloor \dots \left\lfloor \frac{q^{n-k+d/2+1} - 1}{q^{d/2+1} - 1} \left\lfloor \frac{q^{n-k+d/2} - 1}{q^{d/2} - 1} \right\rfloor \dots \right\rfloor \right\rfloor \right\rfloor \right\rfloor$$

 T. Xia and F.-W. Fu. Johnson type bounds on constant dimension codes. *Designs, Codes and Cryptography*, 50:163–172, 2009.

Khaleghi, D. Silva, and F. R. Kschischang. Subspace codes. In *IMA International Conference on Cryptography and Coding*, pages 1–21. Springer, 2009.

T. Etzion and A. Vardy. Error-correcting codes in projective space, *IEEE Transaction on Information Theory*, 2011.

Cyclic subspace codes

$$\mathcal{C} = \cup_V \text{Orb}(V) = \cup_V \{\alpha V : \alpha \in \mathbb{F}_{q^n}^*\}$$

cyclic subspace code

* **Cardinality**

$\mathcal{A}_q(n, d, k) := \max \#$ of codewords a subspace code $\mathcal{C} \subseteq \mathcal{G}_q(n, k)$ may have

* **When $d = 2k - 2$**

Johnson type bound II

$$\mathcal{A}_q(n, 2k - 2, k) \leq \left\lfloor \frac{q^n - 1}{q^k - 1} \left\lfloor \frac{q^{n-1} - 1}{q^{k-1} - 1} \right\rfloor \right\rfloor := J_q(n, 2k - 2, k)$$

Largest known constructions of cyclic subspace codes in $\mathcal{G}_q(rk, k)$ with minimum distance $2k-2$

Johnson type bound II

$$\mathcal{A}_q(rk, 2k-2, k) \leq \left\lfloor \frac{q^{rk}-1}{q^k-1} \left\lfloor \frac{q^{rk-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor := J_q(rk, 2k-2, k) \sim q^{2(r-1)k}$$

as k (or q) $\rightarrow +\infty$

Parameters	Size	Asymptotic behavior
$n = 2k,$ $q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2}q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k-1)(q^{4k}-1)$	$\sim \frac{1}{2}q^{6k}$
$n = rk,$ $r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk}-1),$ $\ell = (\lceil \frac{r}{2} \rceil - 2)$	$\sim q^{(\lceil \frac{r-1}{2} \rceil + r)k}$
$n = rk,$ $r = 2h+1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k-1)^h (q^{rk}-1) + \frac{(q^k-1)^{h-1} (q^{rk}-1)}{q-1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk,$ $r = 2h+1, h \geq 2$	$S_5(rk, k, q) = hq^k (q^k-1)^{h-1} (q^{rk}-1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

Largest known constructions of cyclic subspace codes in $\mathcal{G}_q(rk, k)$ with minimum distance $2k-2$

Johnson type bound II

$$\mathcal{A}_q(rk, 2k-2, k) \leq \left\lfloor \frac{q^{rk}-1}{q^k-1} \left\lfloor \frac{q^{rk-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor := J_q(rk, 2k-2, k) \sim q^{2(r-1)k}$$

as k (or q) $\rightarrow +\infty$

$\mathcal{F}$ family of codes of sizes $S(rk, k, q)$ is
 Asymptotically optimal if

$$\lim_{k(\text{or } q) \rightarrow +\infty} \frac{S(rk, k, q)}{J_q(rk, 2k-2, k)} = 1$$

Parameters	Size	Asymptotic behavior
$n = 2k,$ $q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2}q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k-1)(q^{4k}-1)$	$\sim \frac{1}{2}q^{6k}$
$n = rk,$ $r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk}-1),$ $\ell = (\lceil \frac{r}{2} \rceil - 2)$	$\sim q^{(\lceil \frac{r-1}{2} \rceil + r)k}$
$n = rk,$ $r = 2h+1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k-1)^h (q^{rk}-1) + \frac{(q^k-1)^{h-1} (q^{rk}-1)}{q-1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk,$ $r = 2h+1, h \geq 2$	$S_5(rk, k, q) = hq^k (q^k-1)^{h-1} (q^{rk}-1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

Largest known constructions of cyclic subspace codes in $\mathcal{G}_q(rk, k)$ with minimum distance $2k-2$

R. M. Roth, N. Raviv, & I. Tamo. Construction of Sidon spaces with applications to coding. *IEEE Transactions on Information Theory*, 64(6), 4412-4422, **2018**.

Theorem (Roth, Raviv, Tamo, 2018)

$$n = 2k, q \geq 3, \omega \text{ primitive element of } \mathbb{F}_{q^k}$$

$$\gamma \in \mathbb{F}_{q^n} \setminus \mathbb{F}_{q^k} \text{ s.t. } N_{q^{2k}/q^k}(\gamma) = \omega, \text{ for } h = 1, \dots, \left\lfloor \frac{q-1}{2} \right\rfloor.$$

Let

$${}_2V_{k,h} := \{v + v^q \omega^h \gamma : v \in \mathbb{F}_{q^k}\} \subseteq \mathbb{F}_{q^{2k}} = \mathbb{F}_{q^n}.$$

The code

$${}_2\mathcal{C}_k = \bigcup_{h=1}^{\tau} \text{Orb}({}_2V_{k,h}) \subseteq \mathcal{G}_q(2k, k)$$

$$\text{has minimum distance } 2k - 2 \text{ and size } \left\lfloor \frac{q-1}{2} \right\rfloor \frac{q^{2k} - 1}{q - 1}.$$

Largest known constructions of cyclic subspace codes in $\mathcal{G}_q(rk, k)$ with minimum distance $2k-2$

Parameters	Size	Asymptotic behavior
$n = 2k,$ $q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2}q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k - 1)(q^{4k} - 1)$	$\sim \frac{1}{2}q^{6k}$
$n = rk,$ $r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk} - 1),$ $\ell = (\lceil \frac{r}{2} \rceil - 2)$	$\sim q^{(\lceil \frac{r-1}{2} \rceil + r)k}$
$n = rk,$ $r = 2h + 1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k - 1)^h (q^{rk} - 1) + \frac{(q^k - 1)^{h-1} (q^{rk} - 1)}{q - 1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk,$ $r = 2h + 1, h \geq 2$	$S_5(rk, k, q) = hq^k (q^k - 1)^{h-1} (q^{rk} - 1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

Johnson type bound II

$$\mathcal{A}_q(2k, 2k-2, k) \leq \left\lfloor \frac{q^{2k}-1}{q^k-1} \left\lfloor \frac{q^{2k-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor = J_q(2k, 2k-2, k) \sim q^{2k}$$

as k (or q) $\rightarrow +\infty$

$$\lim_{k(\text{or } q) \rightarrow +\infty} \frac{S(2k, k, q)}{J_q(2k, 2k-2, k)} = \frac{1}{2}$$

R. M. Roth, N. Raviv, & I. Tamo. Construction of Sidon spaces with applications to coding. *IEEE Transactions on Information Theory*, 64(6), 4412-4422, 2018.

Theorem (Roth, Raviv, Tamo, 2018)

$n = 2k, q \geq 3, \omega$ primitive element of $\mathbb{F}_{q^k}$

$\gamma \in \mathbb{F}_{q^n} \setminus \mathbb{F}_{q^k}$ s.t. $N_{q^{2k}/q^k}(\gamma) = \omega$, for $h = 1, \dots, \left\lfloor \frac{q-1}{2} \right\rfloor$.

Let

$${}_2V_{k,h} := \{v + v^q \omega^h \gamma : v \in \mathbb{F}_{q^k}\} \subseteq \mathbb{F}_{q^{2k}} = \mathbb{F}_{q^n}.$$

The code

$${}_2\mathcal{C}_k = \bigcup_{h=1}^{\tau} \text{Orb}({}_2V_{k,h}) \subseteq \mathcal{G}_q(2k, k)$$

has minimum distance $2k-2$ and size $\left\lfloor \frac{q-1}{2} \right\rfloor \frac{q^{2k}-1}{q-1}$.

Largest known constructions of cyclic subspace codes in $\mathcal{G}_q(rk, k)$ with minimum distance $2k-2$

S. Yu, L. Ji. Two new constructions of cyclic subspace codes via Sidon spaces. *Designs, Codes and Cryptography*, 92(11), 3799-3811, **2024**.

Parameters	Size	Asymptotic behavior
$n = 2k,$ $q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2}q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k - 1)(q^{4k} - 1)$	$\sim \frac{1}{2}q^{6k}$
$n = rk,$ $r \geq 3$	$S_3(rk, k, q) = q^{k \frac{q^{\lfloor \frac{r-1}{2} \rfloor} - 1}{q^k - 1}} (q^{rk} - 1),$ $\ell = (\lfloor \frac{r}{2} \rfloor - 2)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk,$ $r = 2h + 1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k - 1)^h (q^{rk} - 1) + \frac{(q^k - 1)^{h-1} (q^{rk} - 1)}{q - 1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk,$ $r = 2h + 1, h \geq 2$	$S_5(rk, k, q) = h q^k (q^k - 1)^{h-1} (q^{rk} - 1) + \frac{q^{rk} - 1}{q^k - 1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

Johnson type bound II

$$\mathcal{A}_q(4k, 2k - 2, k) \leq \left\lfloor \frac{q^{4k} - 1}{q^k - 1} \left\lfloor \frac{q^{4k-1} - 1}{q^{k-1} - 1} \right\rfloor \right\rfloor := J_q(4k, 2k - 2, k) \sim q^{6k}$$

as k (or q) $\rightarrow +\infty$

$$\lim_{k(\text{or } q) \rightarrow +\infty} \frac{S(4k, k, q)}{J_q(4k, 2k - 2, k)} = \frac{1}{2}$$

Largest known constructions of cyclic subspace codes in $\mathcal{G}_q(rk, k)$ with minimum distance $2k-2$

Parameters	Size	Asymptotic behavior
$n = 2k,$ $q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2}q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k - 1)(q^{4k} - 1)$	$\sim \frac{1}{2}q^{6k}$
$n = rk,$ $r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk} - 1),$ $\ell = (\lceil \frac{r}{2} \rceil - 2)$	$\sim q^{(\lceil \frac{r-1}{2} \rceil + r)k}$
$n = rk,$ $r = 2h + 1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k - 1)^h (q^{rk} - 1) + \frac{(q^k - 1)^{h-1} (q^{rk} - 1)}{q - 1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk,$ $r = 2h + 1, h \geq 2$	$S_5(rk, k, q) = h q^k (q^k - 1)^{h-1} (q^{rk} - 1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

Johnson type bound II

$$\mathcal{A}_q(rk, 2k-2, k) \leq \left\lfloor \frac{q^{rk}-1}{q^k-1} \left\lfloor \frac{q^{rk-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor := J_q(rk, 2k-2, k) \sim q^{2(r-1)k}$$

as k (or q) $\rightarrow +\infty$

$$\lim_{k(\text{or } q) \rightarrow +\infty} \frac{S(rk, k, q)}{J_q(rk, 2k-2, k)} = 1 \Leftrightarrow r = 3$$

Zhang, H., Tang, C., & Cao, X. Large optimal cyclic subspace codes. *Discrete Mathematics*, 347(7), 114007, 2024.

Theorem (Zhang, Tang, Cao, 2024)

$n = rk, r \geq 3, \ell := \left\lceil \frac{r}{2} \right\rceil - 2, \omega$ primitive element of $\mathbb{F}_{q^k}$
 $\gamma \in \mathbb{F}_{q^n} \setminus \mathbb{F}_{q^k}$ s.t. $\mathbb{F}_{q^{2k}} = \mathbb{F}_{q^k}(\gamma)$
 $I_{k, \ell_0, \ell} := \{(\alpha_1, \dots, \alpha_{\ell_0}) \in \mathbb{F}_{q^k}^{\ell_0} : \alpha_1, \dots, \alpha_{\ell_0-1} \in \mathbb{F}_{q^k}, \alpha_{\ell_0} \in \mathbb{F}_{q^k}^* \} \times \mathbb{F}_{q^k} \times \{0, 1, \dots, q-2\} \times \{\ell_0, \ell_0+1, \dots, \ell\}$
 For $0 \leq \ell_0 \leq \ell$, define

 ${}_rV_{k, \underline{h}} = \{uP_{k, \underline{\alpha_{\ell_0}}}(\gamma) + \omega^j(u^q + au)\gamma^{\ell_1+1} : u \in \mathbb{F}_{q^k}\} \in \mathcal{G}_q(rk, k)$ for any
 $\underline{h} = (\underline{\alpha_{\ell_0}}, a, j, \ell_1) \in I_{k, \ell_0, \ell}.$

 The code

$${}_r\mathcal{C}_k = \bigcup_{\ell_0=0}^{\ell} \bigcup_{\underline{h} \in I_{k, \ell_0, \ell}} \text{Orb}({}_rV_{k, \underline{h}})$$

 has minimum distance $2k-2$ and size $q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk}-1).$

Take away:

$$\lim_{k(\text{or } q) \rightarrow +\infty} \frac{S(2k, k, q)}{J_q(2k, 2k - 2, k)} = \frac{1}{2} \quad n = 2k$$

R. M. Roth, N. Raviv, & I. Tamo. Construction of Sidon spaces with applications to coding. *IEEE Transactions on Information Theory*, 64(6), 4412-4422, **2018**.

**(Almost) Asymptotically optimal
cyclic subspace codes**

$$\lim_{k(\text{or } q) \rightarrow +\infty} \frac{S(3k, k, q)}{J_q(3k, 2k - 2, k)} = 1 \quad n = 3k$$


Zhang, H., Tang, C., & Cao, X. Large optimal cyclic subspace codes. *Discrete Mathematics*, 347(7), 114007, **2024**.

**Asymptotically optimal
cyclic subspace codes**

Largest known constructions of cyclic subspace codes in $\mathcal{G}_q(rk, k)$ with minimum distance $2k-2$

$$\mathcal{C} = \cup_i \mathcal{C}_i \subseteq \mathcal{G}_q(n, k)$$

$$\mathcal{C}_i \subseteq \mathcal{G}_q(n, k) \quad \forall i$$




Parameters	Size	Asymptotic behavior
$n = 2k,$ $q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2}q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k - 1)(q^{4k} - 1)$	$\sim \frac{1}{2}q^{6k}$
$n = rk,$ $r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk} - 1),$ $\ell = (\lceil \frac{r}{2} \rceil - 2)$	$\sim q^{(\lceil \frac{r-1}{2} \rceil + r)k}$
$n = rk,$ $r = 2h + 1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k - 1)^h (q^{rk} - 1) + \frac{(q^k-1)^{h-1} (q^{rk}-1)}{q-1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk,$ $r = 2h + 1, h \geq 2$	$S_5(rk, k, q) = hq^k (q^k - 1)^{h-1} (q^{rk} - 1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

Common approach

$$\mathcal{C} = \cup_i \mathcal{C}_i \subseteq \mathcal{G}_q(n, k)$$


$$\mathcal{C}_i \subseteq \mathcal{G}_q(n, k) \quad \forall i$$



VS

Our approach

$$\mathcal{C} = \mathcal{C}_t \odot \cdots \odot \mathcal{C}_1 \subseteq \mathcal{G}_q(n, k)$$
$$n = n_t \cdots n_1$$




Our goal

New Constructions of
Asymptotically Optimal cyclic
subspace codes $\mathcal{C} \subseteq \mathcal{G}_q(rk, k)$

$$d = 2k - 2$$

$$|\mathcal{C}| \sim q^{2(r-1)k}$$


$$\mathcal{C} = \mathcal{C}_t \odot \cdots \odot \mathcal{C}_1 \subseteq \mathcal{G}_q(n, k)$$
$$n = n_t \cdots n_1$$



C.C., P. Santonastaso. Asymptotically optimal cyclic
subspace codes, <https://arxiv.org/abs/2507.09290>, **2025**


Nested operation

$$\mathbb{F}_{q^{3k}} = \mathbb{F}_{q^k}(\eta)$$

$$\mathbb{F}_{q^{6k}} = \mathbb{F}_{q^{3k}}(\gamma)$$

$$V_2 = \text{Im}(\Phi) = \{u + u^q\gamma : u \in \mathbb{F}_{q^{3k}}\} \in \mathcal{G}_q(6k, 3k)$$

$$V_1 = \{v + v^q\eta : v \in \mathbb{F}_{q^k}\} \in \mathcal{G}_q(3k, k)$$



$$\Phi : \mathbb{F}_{q^{3k}} \rightarrow \mathbb{F}_{q^{6k}} \quad \text{injective}$$

$$u \mapsto u + u^q\gamma \quad \mathbb{F}_q\text{-linear}$$


Nested operation

$$\mathbb{F}_{q^{3k}} = \mathbb{F}_{q^k}(\eta)$$

$$\mathbb{F}_{q^{6k}} = \mathbb{F}_{q^{3k}}(\gamma)$$

$$V_2 = \text{Im}(\Phi) = \{u + u^q\gamma : u \in \mathbb{F}_{q^{3k}}\} \in \mathcal{G}_q(6k, 3k)$$

$$V_1 = \{v + v^q\eta : v \in \mathbb{F}_{q^k}\} \in \mathcal{G}_q(3k, k)$$



$$\Phi : \mathbb{F}_{q^{3k}} \rightarrow \mathbb{F}_{q^{6k}} \quad \text{injective} \\ u \mapsto u + u^q\gamma \quad \mathbb{F}_q\text{-linear}$$

$$\begin{aligned} V_2 \odot V_1 &:= \{\Phi(u) : u \in V_1\} \\ &= \{u + u^q\gamma : u \in V_1\} \\ &= \{(v + v^q\eta) + (v + v^q\eta)^q\gamma : v \in \mathbb{F}_{q^k}\} \in \mathcal{G}_q(6k, k) \end{aligned}$$


Nested operation

$$V_2 = \text{Im}(\Phi)$$

$$\Phi: \mathbb{F}_{q^m} \rightarrow \mathbb{F}_{q^n}$$

 injective $\mathbb{F}_q$ -linear

$$V_2 \in \mathcal{G}_q(n, m)$$



$$V_2 \odot V_1 := \text{Im}(\Phi|_{V_1}) \in \mathcal{G}_q(n, k)$$

$$V_1 \in \mathcal{G}_q(m, k)$$


Nested operation

$$V_2 = \text{Im}(\Phi)$$

$$\Phi: \mathbb{F}_{q^m} \rightarrow \mathbb{F}_{q^n}$$

 injective $\mathbb{F}_q$ -linear

$$V_2 \in \mathcal{G}_q(n, m)$$



$$V_2 \odot V_1 := \text{Im}(\Phi|_{V_1}) \in \mathcal{G}_q(n, k)$$

$$V_1 \in \mathcal{G}_q(m, k)$$


Nested operation

$$V_2 = \text{Im}(\Phi) \quad \Phi: \mathbb{F}_{q^m} \rightarrow \mathbb{F}_{q^n}$$

injective $\mathbb{F}_q$ -linear

$$V_2 \in \mathcal{G}_q(n, m)$$

$$V_1 \in \mathcal{G}_q(m, k)$$




$$V_2 \odot V_1 := \text{Im}(\Phi|_{V_1}) \in \mathcal{G}_q(n, k)$$

Nested operation

$$V_{2,h_2} = \text{Im}(\Phi_{2,h_2}) \quad \Phi_{2,h_2}: \mathbb{F}_{q^m} \rightarrow \mathbb{F}_{q^n} \\ \forall h_2 = 1, \dots, s_2$$

$$\mathcal{C}_2 = \bigcup_{h_2=1}^{s_2} \text{Orb}(V_{2,h_2}) \subseteq \mathcal{G}_q(n, m)$$

$$\mathcal{C}_1 = \bigcup_{h_1=1}^{s_1} \text{Orb}(V_{1,h_1}) \subseteq \mathcal{G}_q(m, k)$$



$$\mathcal{C}_2 \odot \mathcal{C}_1 := \bigcup_{h_2=1}^{s_2} \bigcup_{h_1=1}^{s_1} \bigcup_{\alpha \in \mathbb{F}_{q^m}^*} \text{Orb}(V_{2,h_2} \odot \alpha V_{1,h_1}) \subseteq \mathcal{G}_q(n, k).$$

Nested operation

$$\mathbb{F}_{q^{3k}} = \mathbb{F}_{q^k}(\gamma_1) \quad V_{1,a_1,j_1} = \{v + \omega_1^{j_1}(v^q + a_1 v)\gamma_1 : v \in \mathbb{F}_{q^k}\} \in \mathcal{G}_q(3k, k)$$

$$\mathbb{F}_{q^{9k}} = \mathbb{F}_{q^{3k}}(\gamma_2) \quad V_{2,a_2,j_2} = \text{Im}(\Phi_{2,a_2,j_2}) = \{u + \omega_2^{j_2}(u^q + u a_2)\gamma_2 : u \in \mathbb{F}_{q^{3k}}\} \in \mathcal{G}_q(9k, 3k)$$

$$\omega_1 \in \mathbb{F}_{q^k} \quad \textbf{Primitive element}$$

$$\omega_2 \in \mathbb{F}_{q^{3k}} \quad \textbf{Primitive element}$$

$$0 \leq j_1, j_2 \leq q-2$$


$$\Phi_{2,a_2,j_2} : \mathbb{F}_{q^{3k}} \rightarrow \mathbb{F}_{q^{9k}}$$

injective $u \mapsto u + \omega_2^{j_2}(u^q + u a_2)\gamma_2$

$\mathbb{F}_q$ -linear

$$\mathcal{C}_2 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_2 \in \mathbb{F}_{q^{3k}}} \text{Orb}(V_{2,a_2,j_2}) \subseteq \mathcal{G}_q(9k, 3k)$$

$$\Phi_{2,a_2}$$




$$\mathcal{C}_2 \odot \mathcal{C}_1 := \bigcup_{0 \leq j_1, j_2 \leq q-2} \bigcup_{a_2 \in \mathbb{F}_{q^{3k}}} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \bigcup_{\alpha \in \mathbb{F}_{q^{3k}}^*} \text{Orb}(V_{2,a_2,j_2} \odot \alpha V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(9k, k)$$

$$\mathcal{C}_1 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \text{Orb}(V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(3k, k)$$

What is the advantage?

Theorem 1 (C.C., P. Santonastaso)

$$\begin{aligned}\mathcal{C}_1 &= \bigcup_{h_1=1}^{s_1} \text{Orb}(V_{1,h_1}) \subseteq \mathcal{G}_q(m, k), & d(\mathcal{C}_1) &= 2k - 2 \\ \mathcal{C}_2 &= \bigcup_{h_2=1}^{s_2} \text{Orb}(V_{2,h_2}) \subseteq \mathcal{G}_q(n, m) & d(\mathcal{C}_2) &= 2m - 2 \\ &V_{2,h_2} = \text{Im}(\Phi_{2,h_2}) \\ &\Phi_{2,h_2}: \mathbb{F}_{q^m} \rightarrow \mathbb{F}_{q^n} \text{ injective } \mathbb{F}_q\text{-linear } \forall h_2 = 1, \dots, s_2\end{aligned}$$



$$\mathcal{C}_2 \odot \mathcal{C}_1 \subseteq \mathcal{G}_q(n, k) \quad d(\mathcal{C}_2 \odot \mathcal{C}_1) = 2k - 2 \quad |\mathcal{C}_2 \odot \mathcal{C}_1| = |\mathcal{C}_2||\mathcal{C}_1|$$

Nested operation

$$\mathbb{F}_{q^{3k}} = \mathbb{F}_{q^k}(\gamma_1) \quad V_{1,a_1,j_1} = \{v + \omega_1^{j_1}(v^q + a_1 v)\gamma_1 : v \in \mathbb{F}_{q^k}\} \in \mathcal{G}_q(3k, k)$$

$$\mathbb{F}_{q^{9k}} = \mathbb{F}_{q^{3k}}(\gamma_2) \quad V_{2,a_2,j_2} = \text{Im}(\Phi_{2,a_2,j_2}) = \{u + \omega_2^{j_2}(u^q + u a_2)\gamma_2 : u \in \mathbb{F}_{q^{3k}}\} \in \mathcal{G}_q(9k, 3k)$$

$\omega_1 \in \mathbb{F}_{q^k}$ **Primitive element**


$\omega_2 \in \mathbb{F}_{q^{3k}}$ **Primitive element**

$$0 \leq j_1, j_2 \leq q - 2$$

$$\Phi_{2,a_2,j_2} : \mathbb{F}_{q^{3k}} \rightarrow \mathbb{F}_{q^{9k}}$$

injective $u \mapsto u + \omega_2^{j_2}(u^q + u a_2)\gamma_2$

$\mathbb{F}_q$ -linear



$$\mathcal{C}_1 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \text{Orb}(V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(3k, k)$$

$$\mathcal{C}_2 = \bigcup_{0 \leq j_2 \leq q-2} \bigcup_{a_2 \in \mathbb{F}_{q^{3k}}} \text{Orb}(V_{2,a_2,j_2}) \subseteq \mathcal{G}_q(9k, 3k)$$

$$\mathcal{C}_2 \odot \mathcal{C}_1 := \bigcup_{0 \leq j_1, j_2 \leq q-2} \bigcup_{a_2 \in \mathbb{F}_{q^{3k}}} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \bigcup_{\alpha \in \mathbb{F}_{q^{3k}}^*} \text{Orb}(V_{2,a_2,j_2} \odot \alpha V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(9k, k)$$

$$d(\mathcal{C}_2 \odot \mathcal{C}_1) = 2k - 2$$

Nested operation

$$\mathbb{F}_{q^{3k}} = \mathbb{F}_{q^k}(\gamma_1) \quad V_{1,a_1,j_1} = \{v + \omega_1^{j_1}(v^q + a_1 v)\gamma_1 : v \in \mathbb{F}_{q^k}\} \in \mathcal{G}_q(3k, k)$$

$$\mathbb{F}_{q^{9k}} = \mathbb{F}_{q^{3k}}(\gamma_2) \quad V_{2,a_2,j_2} = \text{Im}(\Phi_{2,a_2,j_2}) = \{u + \omega_2^{j_2}(u^q + u a_2)\gamma_2 : u \in \mathbb{F}_{q^{3k}}\} \in \mathcal{G}_q(9k, 3k)$$

$$\omega_1 \in \mathbb{F}_{q^k} \text{ Primitive element}$$


$$\omega_2 \in \mathbb{F}_{q^{3k}} \text{ Primitive element}$$

$$0 \leq j_1, j_2 \leq q - 2$$

$$\Phi_{2,a_2,j_2} : \mathbb{F}_{q^{3k}} \rightarrow \mathbb{F}_{q^{9k}}$$

$$\text{injective} \quad u \mapsto u + \omega_2^{j_2}(u^q + u a_2)\gamma_2$$

$$\mathbb{F}_q\text{-linear}$$



$$\mathcal{C}_1 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \text{Orb}(V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(3k, k)$$

$$\mathcal{C}_2 = \bigcup_{0 \leq j_2 \leq q-2} \bigcup_{a_2 \in \mathbb{F}_{q^{3k}}} \text{Orb}(V_{2,a_2,j_2}) \subseteq \mathcal{G}_q(9k, 3k)$$

$$\mathcal{C}_2 \odot \mathcal{C}_1 := \bigcup_{0 \leq j_1, j_2 \leq q-2} \bigcup_{a_2 \in \mathbb{F}_{q^{3k}}} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \bigcup_{\alpha \in \mathbb{F}_{q^{3k}}^*} \text{Orb}(V_{2,a_2,j_2} \odot \alpha V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(9k, k)$$

$$d(\mathcal{C}_2 \odot \mathcal{C}_1) = 2k - 2$$

$$|\mathcal{C}_2 \odot \mathcal{C}_1| = |\mathcal{C}_2| |\mathcal{C}_1| = q^{4k}(q^{3k} - 1)(q^{9k} - 1) \sim q^{16k}$$

Nested operation


$$\mathbb{F}_{q^{3k}} = \mathbb{F}_{q^k}(\gamma_1) \qquad V_{1,a_1,j_1} = \{v + \omega_1^{j_1}(v^q + a_1v)\gamma_1 : v \in \mathbb{F}_{q^k}\} \in \mathcal{G}_q(3k,k)$$

$$\mathbb{F}_{q^{9k}} = \mathbb{F}_{q^{3k}}(\gamma_2) \qquad V_{2,a_2,j_2} = \text{Im}(\Phi_{2,a_2,j_2}) = \{u + \omega_2^{j_2}(u^q + ua_2)\gamma_2 : u \in \mathbb{F}_{q^{3k}}\} \in \mathcal{G}_q(9k,3k)$$

$$\omega_1 \in \mathbb{F}_{q^k} \text{ Primitive element} \qquad \Phi_{2,a_2,j_2} : \mathbb{F}_{q^{3k}} \rightarrow \mathbb{F}_{q^{9k}}$$

$$\omega_2 \in \mathbb{F}_{q^{3k}} \text{ Primitive element} \qquad \text{injective} \quad u \mapsto u + \omega_2^{j_2}(u^q + ua_2)\gamma_2$$

$$0 \leq j_1, j_2 \leq q-2 \qquad \mathbb{F}_q\text{-linear}$$



$$\mathcal{C}_1 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \text{Orb}(V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(3k,k)$$

$$\mathcal{C}_2 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_2 \in \mathbb{F}_{q^{3k}}} \text{Orb}(V_{2,a_2,j_2}) \subseteq \mathcal{G}_q(9k,3k)$$

$$\mathcal{C}_2 \odot \mathcal{C}_1 := \bigcup_{0 \leq j_1, j_2 \leq q-2} \bigcup_{a_2 \in \mathbb{F}_{q^{3k}}} \bigcup_{a_1 \in \mathbb{F}_{q^k}}$$

Johnson type bound II

$$\mathcal{A}_q(9k,2k-2,k) \leq \left\lfloor \frac{q^{9k}-1}{q^k-1} \left\lfloor \frac{q^{9k-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor := J_q(9k,2k-2,k) \sim q^{16k}$$

as k (or q) $\rightarrow +\infty$

$$d(\mathcal{C}_2 \odot \mathcal{C}_1) = 2k-2$$

$$|\mathcal{C}_2 \odot \mathcal{C}_1| = |\mathcal{C}_2||\mathcal{C}_1| = q^{4k}(q^{3k}-1)(q^{9k}-1) \sim q^{16k}$$



Remark

$$\mathbb{F}_{q^{3k}} = \mathbb{F}_{q^k}(\gamma_1)$$

$$V_{1,a_1,j_1} = \{v + \omega_1^{j_1}(v^q + a_1v)\gamma_1 : v \in \mathbb{F}_{q^k}\} \in \mathcal{G}_q(3k,k)$$

$$\mathbb{F}_{q^{9k}} = \mathbb{F}_{q^{3k}}(\gamma_2)$$

$$V_{2,a_2,j_2} = \text{Im}(\Phi_{2,a_2,j_2}) = \{u + \omega_2^{j_2}(u^q + ua_2)\gamma_2 : u \in \mathbb{F}_{q^{3k}}\} \in \mathcal{G}_q(9k,3k)$$

$$\omega_1 \in \mathbb{F}_{q^k} \text{ Primitive}$$

$$\omega_2 \in \mathbb{F}_{q^{3k}}$$

$$0 <$$

$$n = 9k = 3^2k$$

$$\begin{aligned} &\text{injective} \\ &\mathbb{F}_q\text{-linear} \\ &\Phi_{2,a_2,j_2} : \mathbb{F}_{q^{3k}} \rightarrow \mathbb{F}_{q^{9k}} \\ &u \mapsto u + \omega_2^{j_2}(u^q + ua_2)\gamma_2 \end{aligned}$$

$$\mathcal{C}_1 = \cup_{0 \leq j_1 \leq q-2} \cup_{a_1 \in \mathbb{F}_{q^k}} \text{Orb}(V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(3k,k)$$

$$\mathcal{C}_2 = \cup_{0 \leq j_1 \leq q-2} \cup_{a_2 \in \mathbb{F}_{q^{3k}}} \text{Orb}(V_{2,a_2,j_2}) \subseteq \mathcal{G}_q(9k,3k)$$

Johnson type bound II

$$\mathcal{A}_q(rk,2k-2,k) \leq \left\lfloor \frac{q^{rk}-1}{q^k-1} \left\lfloor \frac{q^{rk-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor \sim q^{2(r-1)k}$$

as k (or q) $\rightarrow +\infty$

$$\lim_{k(\text{or } q) \rightarrow +\infty} \frac{S(rk,k,q)}{J_q(rk,2k-2,k)} = 1 \Leftrightarrow r = 3$$

Parameters	Size	Asymptotic behavior
$n = 2k, q > 2$	$S_1(2k,k,q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2}q^{2k}$
$n = 4k$	$S_2(4k,k,q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k-1)(q^{4k}-1)$	$\sim \frac{1}{2}q^{6k}$
$n = rk, r \geq 3$	$S_3(rk,k,q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk}-1), \ell = \lfloor \frac{r}{2} \rfloor - 2$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_4(rk,k,q) = h \left((q^k-1)^h (q^{rk}-1) + \frac{(q^k-1)^{h-1} (q^{rk}-1)}{q-1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_5(rk,k,q) = hq^k (q^k-1)^{h-1} (q^{rk}-1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

Johnson type bound II

$$\mathcal{A}_q(9k,2k-2,k) \leq \left\lfloor \frac{q^{9k}-1}{q^k-1} \left\lfloor \frac{q^{9k-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor := J_q(9k,2k-2,k) \sim q^{16k}$$

as k (or q) $\rightarrow +\infty$

$\mathcal{C} \subseteq \mathcal{G}_q(n',k')$ asymptotically optimal
when $n'/k' = 3$

$$d(\mathcal{C}_2 \odot \mathcal{C}_1) = 2k - 2$$

$$|\mathcal{C}_2 \odot \mathcal{C}_1| = |\mathcal{C}_2||\mathcal{C}_1| = q^{4k}(q^{3k}-1)(q^{9k}-1) \sim q^{16k}$$

Strategy

$$n = 6k = 2 \cdot 3k$$

$$\mathcal{C}_2 = \text{Orb}(V_2) = \{\beta V_2 : \beta \in \mathbb{F}_{q^{6k}}\} \subseteq \mathcal{G}_q(6k, 3k)$$

Johnson type bound II

$$\mathcal{A}_q(2k, 2k-2, k) \leq \left\lfloor \frac{q^{2k}-1}{q^k-1} \left\lfloor \frac{q^{2k-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor \sim q^{2k}$$

as k (or q) $\rightarrow +\infty$

Parameters	Size	Asymptotic behavior
$n = 2k, q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2} q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k-1)(q^{4k}-1)$	$\sim \frac{1}{2} q^{6k}$
$n = rk, r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk}-1), \ell = \lfloor \frac{r}{2} \rfloor - 2$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k-1)^h (q^{rk}-1) + \frac{(q^k-1)^{h-1} (q^{rk}-1)}{q-1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_5(rk, k, q) = h q^k (q^k-1)^{h-1} (q^{rk}-1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

$\mathcal{C} \subseteq \mathcal{G}_q(n', k')$ asymptotically optimal
(within a factor of $1/2 + o_k(1)$)
when $n'/k' = 2$

$$\mathcal{C}_1 = \text{Orb}(V_1) = \{\alpha V_1 : \alpha \in \mathbb{F}_{q^{3k}}\} \subseteq \mathcal{G}_q(3k, k)$$

Johnson type bound II

$$\mathcal{A}_q(rk, 2k-2, k) \leq \left\lfloor \frac{q^{rk}-1}{q^k-1} \left\lfloor \frac{q^{rk-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor \sim q^{2(r-1)k}$$

as k (or q) $\rightarrow +\infty$

Parameters	Size	Asymptotic behavior
$n = 2k, q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2} q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k-1)(q^{4k}-1)$	$\sim \frac{1}{2} q^{6k}$
$n = rk, r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk}-1), \ell = \lfloor \frac{r}{2} \rfloor - 2$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k-1)^h (q^{rk}-1) + \frac{(q^k-1)^{h-1} (q^{rk}-1)}{q-1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_5(rk, k, q) = h q^k (q^k-1)^{h-1} (q^{rk}-1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

$$\lim_{k(\text{or } q) \rightarrow +\infty} \frac{S(rk, k, q)}{J_q(rk, 2k-2, k)} = 1 \Leftrightarrow r = 3$$

$\mathcal{C} \subseteq \mathcal{G}_q(n', k')$ asymptotically optimal
when $n'/k' = 3$

Strategy

$$n = 6k = 2 \cdot 3k$$

$$\mathcal{C}_2 = \text{Orb}(V_2) = \{\beta V_2 : \beta \in \mathbb{F}_{q^{6k}}\} \subseteq \mathcal{G}_q(6k, 3k)$$

Johnson type bound II

$$\mathcal{A}_q(2k, 2k-2, k) \leq \left\lfloor \frac{q^{2k}-1}{q^k-1} \left\lfloor \frac{q^{2k-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor \sim q^{2k} \quad \text{as } k \text{ (or } q) \rightarrow +\infty$$

Parameters	Size	Asymptotic behavior
$n = 2k, q > 2$	$S_1(2k, k, q) = \left\lfloor \frac{q-1}{2} \right\rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2} q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \left\lfloor \frac{q^k-2}{2} \right\rfloor (q^k-1)(q^{4k}-1)$	$\sim \frac{1}{2} q^{6k}$
$n = rk, r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk}-1), \ell = \left\lceil \frac{r}{2} \right\rceil - 2$	$\sim q^{(\lceil \frac{r-1}{2} \rceil + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k-1)^h (q^{rk}-1) + \frac{(q^k-1)^{h-1} (q^{rk}-1)}{q-1} \right)$	$\sim q^{(\lceil \frac{r-1}{2} \rceil + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_5(rk, k, q) = h q^k (q^k-1)^{h-1} (q^{rk}-1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lceil \frac{r-1}{2} \rceil + r)k}$

$\mathcal{C} \subseteq \mathcal{G}_q(n', k')$ asymptotically optimal
(within a factor of $1/2 + o_k(1)$)
when $n'/k' = 2$

$$\mathcal{C}_1 = \text{Orb}(V_1) = \{\alpha V_1 : \alpha \in \mathbb{F}_{q^{3k}}\} \subseteq \mathcal{G}_q(3k, k)$$

Johnson type bound II

$$\mathcal{A}_q(rk, 2k-2, k) \leq \left\lfloor \frac{q^{rk}-1}{q^k-1} \left\lfloor \frac{q^{rk-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor \sim q^{2(r-1)k} \quad \text{as } k \text{ (or } q) \rightarrow +\infty$$

Parameters	Size	Asymptotic behavior
$n = 2k, q > 2$	$S_1(2k, k, q) = \left\lfloor \frac{q-1}{2} \right\rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2} q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \left\lfloor \frac{q^k-2}{2} \right\rfloor (q^k-1)(q^{4k}-1)$	$\sim \frac{1}{2} q^{6k}$
$n = rk, r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk}-1), \ell = \left\lceil \frac{r}{2} \right\rceil - 2$	$\sim q^{(\lceil \frac{r-1}{2} \rceil + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k-1)^h (q^{rk}-1) + \frac{(q^k-1)^{h-1} (q^{rk}-1)}{q-1} \right)$	$\sim q^{(\lceil \frac{r-1}{2} \rceil + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_5(rk, k, q) = h q^k (q^k-1)^{h-1} (q^{rk}-1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lceil \frac{r-1}{2} \rceil + r)k}$

$\lim_{k(\text{or } q) \rightarrow +\infty} \frac{S(rk, k, q)}{J_q(rk, 2k-2, k)} = 1 \Leftrightarrow r = 3$

$\mathcal{C} \subseteq \mathcal{G}_q(n', k')$ asymptotically optimal
when $n'/k' = 3$

Asymptotically optimal cyclic subspace codes

$$\mathbb{F}_{q^{3k}} = \mathbb{F}_{q^k}(\gamma_1)$$

$$V_{1,a_1,j_1} = \{v + \omega_1^{j_1}(v^q + a_1v)\gamma_1 : v \in \mathbb{F}_{q^k}\} \in \mathcal{G}_q(3k, k)$$

$$\mathbb{F}_{q^{6k}} = \mathbb{F}_{q^{3k}}(\gamma)$$


$$V_2 = \text{Im}(\Phi) = \{u + u^q\gamma : u \in \mathbb{F}_{q^{3k}}\} \in \mathcal{G}_q(6k, 3k)$$

injective

$$\Phi: \mathbb{F}_{q^{3k}} \rightarrow \mathbb{F}_{q^{6k}} \\ u \mapsto u + u^q\gamma$$

$\omega_1 \in \mathbb{F}_{q^k}$ **Primitive element**

$$0 \leq j_1 \leq q-2$$



$$\mathcal{C}_1 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \text{Orb}(V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(3k, k)$$

$$\mathcal{C}_2 = \text{Orb}(V_2) = \{\beta V_2 : \beta \in \mathbb{F}_{q^{6k}}\} \subseteq \mathcal{G}_q(6k, 3k)$$

$$\mathcal{C}_2 \odot \mathcal{C}_1 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \bigcup_{\alpha \in \mathbb{F}_{q^{3k}}^*} \text{Orb}(V_2 \odot \alpha V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(6k, k)$$

Asymptotically optimal cyclic subspace codes

$$\mathbb{F}_{q^{3k}} = \mathbb{F}_{q^k}(\gamma_1)$$

$$V_{1,a_1,j_1} = \{v + \omega_1^{j_1}(v^q + a_1 v)\gamma_1 : v \in \mathbb{F}_{q^k}\} \in \mathcal{G}_q(3k, k)$$

$$\mathbb{F}_{q^{6k}} = \mathbb{F}_{q^{3k}}(\gamma)$$


$$V_2 = \text{Im}(\Phi) = \{u + u^q \gamma : u \in \mathbb{F}_{q^{3k}}\} \in \mathcal{G}_q(6k, 3k)$$

injective

$$\Phi: \mathbb{F}_{q^{3k}} \rightarrow \mathbb{F}_{q^{6k}} \\ u \mapsto u + u^q \gamma$$

$\omega_1 \in \mathbb{F}_{q^k}$ **Primitive element**

$$0 \leq j_1 \leq q-2$$



$$\mathcal{C}_1 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \text{Orb}(V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(3k, k)$$

$$\mathcal{C}_2 = \text{Orb}(V_2) = \{\beta V_2 : \beta \in \mathbb{F}_{q^{6k}}\} \subseteq \mathcal{G}_q(6k, 3k)$$

$$\mathcal{C}_2 \odot \mathcal{C}_1 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \bigcup_{\alpha \in \mathbb{F}_{q^{3k}}^*} \text{Orb}(V_2 \odot \alpha V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(6k, k)$$

$$d(\mathcal{C}_2 \odot \mathcal{C}_1) = 2k - 2$$

Asymptotically optimal cyclic subspace codes

$$\mathbb{F}_{q^{3k}} = \mathbb{F}_{q^k}(\gamma_1)$$

$$V_{1,a_1,j_1} = \{v + \omega_1^{j_1}(v^q + a_1 v)\gamma_1 : v \in \mathbb{F}_{q^k}\} \in \mathcal{G}_q(3k, k)$$

$$\mathbb{F}_{q^{6k}} = \mathbb{F}_{q^{3k}}(\gamma)$$


$$V_2 = \text{Im}(\Phi) = \{u + u^q \gamma : u \in \mathbb{F}_{q^{3k}}\} \in \mathcal{G}_q(6k, 3k)$$

injective

$$\Phi: \mathbb{F}_{q^{3k}} \rightarrow \mathbb{F}_{q^{6k}} \\ u \mapsto u + u^q \gamma$$

$\omega_1 \in \mathbb{F}_{q^k}$ **Primitive element**

$$0 \leq j_1 \leq q-2$$



$$\mathcal{C}_1 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \text{Orb}(V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(3k, k)$$

$$\mathcal{C}_2 = \text{Orb}(V_2) = \{\beta V_2 : \beta \in \mathbb{F}_{q^{6k}}\} \subseteq \mathcal{G}_q(6k, 3k)$$

$$\mathcal{C}_2 \odot \mathcal{C}_1 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \bigcup_{\alpha \in \mathbb{F}_{q^{3k}}^*} \text{Orb}(V_2 \odot \alpha V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(6k, k)$$

$$d(\mathcal{C}_2 \odot \mathcal{C}_1) = 2k - 2$$

$$|\mathcal{C}_2 \odot \mathcal{C}_1| = |\mathcal{C}_2| |\mathcal{C}_1| = \left\lfloor \frac{q-1}{2} \right\rfloor \frac{q^{6k}-1}{q-1} q^k (q^{3k}-1) \sim \frac{1}{2} q^{10k}$$

Asymptotically optimal cyclic subspace codes

$$\mathbb{F}_{q^{3k}} = \mathbb{F}_{q^k}(\gamma_1)$$

$$V_{1,a_1,j_1} = \{v + \omega_1^{j_1}(v^q + a_1 v)\gamma_1 : v \in \mathbb{F}_{q^k}\} \in \mathcal{G}_q(3k, k)$$


$$\mathbb{F}_{q^{6k}} = \mathbb{F}_{q^{3k}}(\gamma)$$

$$V_2 = \text{Im}(\Phi) = \{u + u^q\gamma : u \in \mathbb{F}_{q^{3k}}\} \in \mathcal{G}_q(6k, 3k)$$

$$\begin{aligned} \Phi: \mathbb{F}_{q^{3k}} &\rightarrow \mathbb{F}_{q^{6k}} \\ u &\mapsto u + u^q\gamma \\ \text{injective} \\ \mathbb{F}_q\text{-linear} \end{aligned}$$

$\omega_1 \in \mathbb{F}_{q^k}$ **Primitive element**

$$0 \leq j_1 \leq q-2$$



$$\mathcal{C}_1 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \text{Orb}(V_{1,a_1,j_1}) \subseteq \mathcal{G}_q(3k, k)$$

$$\mathcal{C}_2 = \text{Orb}(V_2) = \{\beta V_2 : \beta \in \mathbb{F}_{q^{6k}}\} \subseteq \mathcal{G}_q(6k, 3k)$$

$$\mathcal{C}_2 \odot \mathcal{C}_1 = \bigcup_{0 \leq j_1 \leq q-2} \bigcup_{a_1 \in \mathbb{F}_{q^k}} \bigcup_{\alpha \in \mathbb{F}_{q^{3k}}^*}$$

$$d(\mathcal{C}_2 \odot \mathcal{C}_1) =$$

Johnson type bound II

$$\mathcal{A}_q(6k, 2k-2, k) \leq \left\lfloor \frac{q^{6k}-1}{q^k-1} \left\lfloor \frac{q^{6k-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor := J_q(6k, 2k-2, k) \sim q^{10k}$$

as k (or q) $\rightarrow +\infty$

$$|\mathcal{C}_2 \odot \mathcal{C}_1| = |\mathcal{C}_2| |\mathcal{C}_1| = \left\lfloor \frac{q-1}{2} \right\rfloor \frac{q^{6k}-1}{q-1} q^k (q^{3k}-1) \sim \frac{1}{2} q^{10k}$$

Generalizing...

Theorem 2 (C.C., P. Santonastaso)


$$\mathcal{C}_1 = \bigcup_{h_1=1}^{s_1} \text{Orb}(V_{1,h_1}) = \{U_{1,1}, \dots, U_{1,t_1}\} \subseteq \mathcal{G}_q(m, k)$$

$$\mathcal{C}_i = \bigcup_{h_i=1}^{s_i} \text{Orb}(V_{i,h_i}) = \{U_{i,1}, \dots, U_{i,t_i}\} \subseteq \mathcal{G}_q(r_{i-1} \cdots r_1 m, r_{i-2} \cdots r_1 m)$$

$$n = r_{e-1} \cdots r_1 m, e \geq 2$$

$$d(\mathcal{C}_1) = 2k - 2$$

$$d(\mathcal{C}_i) = 2(r_{i-2} \cdots r_1 m) - 2 \quad \forall i = 2, \dots, e, e \geq 2$$



$$\mathcal{C}_e \odot \cdots \odot \mathcal{C}_1 \subseteq \mathcal{G}_q(n, k)$$

$$d(\mathcal{C}_e \odot \cdots \odot \mathcal{C}_1) = 2k - 2 \quad |\mathcal{C}_e \odot \cdots \odot \mathcal{C}_1| = |\mathcal{C}_e| \cdots |\mathcal{C}_1|$$

Consequences

$$n = 3^e k$$

$$\mathcal{C}_e \odot \cdots \odot \mathcal{C}_1 \subseteq \mathcal{G}_q(3^e k, k)$$

$$\mathcal{C}_1 \subseteq \mathcal{G}_q(3k, k)$$

$$\mathcal{C}_2 \subseteq \mathcal{G}_q(3^2 k, 3k)$$

⋮

$$\mathcal{C}_e \subseteq \mathcal{G}_q(3^e k, 3^{e-1} k)$$

Johnson type bound II

$$\mathcal{A}_q(rk, 2k-2, k) \leq \left\lfloor \frac{q^{rk}-1}{q^k-1} \left\lfloor \frac{q^{rk-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor \sim q^{2(r-1)k}$$

as k (or q) $\rightarrow +\infty$

$$\lim_{k(\text{or } q) \rightarrow +\infty} \frac{S(rk, k, q)}{J_q(rk, 2k-2, k)} = 1 \Leftrightarrow r = 3$$

Parameters	Size	Asymptotic behavior
$n = 2k,$ $q > 2$	$S_1(2k, k, q) = \left\lfloor \frac{q-1}{2} \right\rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2} q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \left\lfloor \frac{q^k-2}{2} \right\rfloor (q^k-1)(q^{4k}-1)$	$\sim \frac{1}{2} q^{6k}$
$n = rk,$ $r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(r+1)k}-1}{q^k-1} (q^{rk}-1),$ $\ell = \left(\left\lfloor \frac{r}{2} \right\rfloor - 2\right)$	$\sim q^{(\lceil \frac{r-1}{2} \rceil + r)k}$
$n = rk,$ $r = 2h+1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k-1)^h (q^{rk}-1) + \frac{(q^k-1)^{h-1} (q^{rk}-1)}{q-1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk,$ $r = 2h+1, h \geq 2$	$S_5(rk, k, q) = h q^k (q^k-1)^{h-1} (q^{rk}-1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

$\mathcal{C} \subseteq \mathcal{G}_q(n', k')$ asymptotically optimal
when $n'/k' = 3$

Consequences

$$n = 3^e k$$

$$\mathcal{C}_e \odot \cdots \odot \mathcal{C}_1 \subseteq \mathcal{G}_q(3^e k, k)$$

$$\mathcal{C}_1 \subseteq \mathcal{G}_q(3k, k)$$

$$\mathcal{C}_2 \subseteq \mathcal{G}_q(3^2 k, 3k)$$

⋮

$$\mathcal{C}_e \subseteq \mathcal{G}_q(3^e k, 3^{e-1} k)$$

$$|\mathcal{C}_e \odot \cdots \odot \mathcal{C}_1| \sim q^{2(3^e - 1)k}$$



Johnson type bound II

$$\mathcal{A}_q(rk, 2k - 2, k) \leq \left\lfloor \frac{q^{rk} - 1}{q^k - 1} \left\lfloor \frac{q^{rk-1} - 1}{q^{k-1} - 1} \right\rfloor \right\rfloor \sim q^{2(r-1)k}$$

as k (or q) $\rightarrow +\infty$

$\lim_{k(\text{or } q) \rightarrow +\infty} \frac{S(rk, k, q)}{J_q(rk, 2k - 2, k)} = 1 \Leftrightarrow r = 3$

Parameters	Size	Asymptotic behavior
$n = 2k,$ $q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2}q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k - 1)(q^{4k} - 1)$	$\sim \frac{1}{2}q^{6k}$
$n = rk,$ $r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk} - 1),$ $\ell = (\lfloor \frac{r}{2} \rfloor - 2)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk,$ $r = 2h + 1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k - 1)^h (q^{rk} - 1) + \frac{(q^k - 1)^{h-1} (q^{rk} - 1)}{q - 1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk,$ $r = 2h + 1, h \geq 2$	$S_5(rk, k, q) = h q^k (q^k - 1)^{h-1} (q^{rk} - 1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

Johnson type bound II

$$\mathcal{A}_q(3^e k, 2k - 2, k) \leq \left\lfloor \frac{q^{3^e k} - 1}{q^k - 1} \left\lfloor \frac{q^{3^e k-1} - 1}{q^{k-1} - 1} \right\rfloor \right\rfloor := J_q(3^e k, 2k - 2, k) \sim q^{2(3^e - 1)k}$$

as k (or q) $\rightarrow +\infty$

Consequences

$$n = 2^e 3^\ell k, e \geq 1, \ell \geq 0$$

$$\mathcal{C}_{e+\ell} \odot \dots \odot \mathcal{C}_1 \subseteq \mathcal{G}_q(2^e 3^\ell k, k)$$

$$\mathcal{C}_1 \subseteq \mathcal{G}_q(3k, k)$$

$$\mathcal{C}_2 \subseteq \mathcal{G}_q(3^2 k, 3k)$$

⋮

$$\mathcal{C}_{e+\ell} \subseteq \mathcal{G}_q(2^\ell 3^e k, 2^{\ell-1} 3^e k)$$

Johnson type bound II

$$\mathcal{A}_q(2k, 2k-2, k) \leq \left\lfloor \frac{q^{2k}-1}{q^k-1} \left\lfloor \frac{q^{2k-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor \sim q^{2k}$$

as k (or q) $\rightarrow +\infty$

Parameters	Size	Asymptotic behavior
$n = 2k, q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2} q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k-1)(q^{4k}-1)$	$\sim \frac{1}{2} q^{6k}$
$n = rk, r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk}-1), \ell = (\lfloor \frac{r}{2} \rfloor - 2)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k-1)^h (q^{rk}-1) + \frac{(q^k-1)^{h-1} (q^{rk}-1)}{q-1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_5(rk, k, q) = h q^k (q^k-1)^{h-1} (q^{rk}-1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

$\mathcal{C} \subseteq \mathcal{G}_q(n', k')$ asymptotically optimal
(within a factor of $1/2 + o_k(1)$)
when $n'/k' = 2$

$\mathcal{C} \subseteq \mathcal{G}_q(n', k')$ asymptotically optimal
when $n'/k' = 3$

Johnson type bound II

$$\mathcal{A}_q(rk, 2k-2, k) \leq \left\lfloor \frac{q^{rk}-1}{q^k-1} \left\lfloor \frac{q^{rk-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor \sim q^{2(r-1)k}$$

as k (or q) $\rightarrow +\infty$

Parameters	Size	Asymptotic behavior
$n = 2k, q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2} q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k-1)(q^{4k}-1)$	$\sim \frac{1}{2} q^{6k}$
$n = rk, r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk}-1), \ell = (\lfloor \frac{r}{2} \rfloor - 2)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k-1)^h (q^{rk}-1) + \frac{(q^k-1)^{h-1} (q^{rk}-1)}{q-1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_5(rk, k, q) = h q^k (q^k-1)^{h-1} (q^{rk}-1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

$$\lim_{k(\text{or } q) \rightarrow +\infty} \frac{S(rk, k, q)}{J_q(rk, 2k-2, k)} = 1 \Leftrightarrow r = 3$$

Consequences

$$n = 2^e 3^\ell k, e \geq 1, \ell \geq 0$$

$$\mathcal{C}_{e+\ell} \odot \cdots \odot \mathcal{C}_1 \subseteq \mathcal{G}_q(2^e 3^\ell k, k)$$

$$\mathcal{C}_1 \subseteq \mathcal{G}_q(3k, k)$$

$$\mathcal{C}_2 \subseteq \mathcal{G}_q(3^2 k, 3k)$$

⋮

$$\mathcal{C}_{e+\ell} \subseteq \mathcal{G}_q(2^\ell 3^e k, 2^{\ell-1} 3^e k)$$

$$|\mathcal{C}_{e+\ell} \odot \cdots \odot \mathcal{C}_1| \sim \frac{1}{2^e} q^{2(3^\ell 2^e - 1)k}$$



Johnson type bound II

$$\mathcal{A}_q(2k, 2k-2, k) \leq \left\lfloor \frac{q^{2k}-1}{q^k-1} \left\lfloor \frac{q^{2k-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor \sim q^{2k}$$

as k (or q) $\rightarrow +\infty$

$\rightarrow$

Parameters	Size	Asymptotic behavior
$n = 2k, q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2} q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k-1)(q^{4k}-1)$	$\sim \frac{1}{2} q^{6k}$
$n = rk, r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk}-1), \ell = (\lfloor \frac{r}{2} \rfloor - 2)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k-1)^h (q^{rk}-1) + \frac{(q^k-1)^{h-1} (q^{rk}-1)}{q-1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_5(rk, k, q) = h q^k (q^k-1)^{h-1} (q^{rk}-1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

$\mathcal{C} \subseteq \mathcal{G}_q(n', k')$ asymptotically optimal
(within a factor of $1/2 + o_k(1)$)
when $n'/k' = 2$

Johnson type bound II

$$\mathcal{A}_q(3^e k, 2k-2, k) \leq \left\lfloor \frac{q^{2^e 3^\ell k}-1}{q^k-1} \left\lfloor \frac{q^{2^e 3^\ell k-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor := J_q(2^e 3^\ell k, 2k-2, k) \sim q^{2(2^e 3^\ell - 1)k}$$

as k (or q) $\rightarrow +\infty$

otically optimal
 $= 3$

Johnson type bound II

$$\mathcal{A}_q(rk, 2k-2, k) \leq \left\lfloor \frac{q^{rk}-1}{q^k-1} \left\lfloor \frac{q^{rk-1}-1}{q^{k-1}-1} \right\rfloor \right\rfloor \sim q^{2(r-1)k}$$

as k (or q) $\rightarrow +\infty$

$\rightarrow$

Parameters	Size	Asymptotic behavior
$n = 2k, q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2} q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k-1)(q^{4k}-1)$	$\sim \frac{1}{2} q^{6k}$
$n = rk, r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk}-1), \ell = (\lfloor \frac{r}{2} \rfloor - 2)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_4(rk, k, q) = h \left((q^k-1)^h (q^{rk}-1) + \frac{(q^k-1)^{h-1} (q^{rk}-1)}{q-1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk, r = 2h+1, h \geq 2$	$S_5(rk, k, q) = h q^k (q^k-1)^{h-1} (q^{rk}-1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

$\lim_{k(\text{or } q) \rightarrow +\infty} \frac{S(rk, k, q)}{J_q(rk, 2k-2, k)} = 1 \Leftrightarrow r = 3$

General construction

Theorem 3 (C.C., P. Santonastaso)

$$n = 2^e p_z^{e_z} \cdots p_1^{e_1} k \quad 2 < p_z < \cdots < p_1 \quad e, e_1, \dots, e_z \geq 0$$

For k (or q) sufficiently large

$$|\mathcal{C}_{e+e_z+\dots+e_1} \odot \cdots \odot \mathcal{C}_1| \sim \frac{1}{2^e} q^{2 \prod_{i=1}^e p_i^{e_i} k (2^e - 1)} q^{\sum_{i=1}^z \left(\frac{(3p_i - 1)(p_i^{e_i} - 1)}{2(p_i - 1)} \prod_{j=1}^{i-1} p_j^{e_j} k \right)} >$$

Parameters	Size	Asymptotic behavior
$n = 2k,$ $q > 2$	$S_1(2k, k, q) = \lfloor \frac{q-1}{2} \rfloor \frac{q^{2k}-1}{q-1}$	$\sim \frac{1}{2} q^{2k}$
$n = 4k$	$S_2(4k, k, q) = \lfloor \frac{q^k-2}{2} \rfloor (q^k - 1)(q^{4k} - 1)$	$\sim \frac{1}{2} q^{6k}$
$n = rk,$ $r \geq 3$	$S_3(rk, k, q) = q^k \frac{q^{(\ell+1)k}-1}{q^k-1} (q^{rk} - 1),$ $\ell = (\lceil \frac{r}{2} \rceil - 2)$	$\sim q^{(\lceil \frac{r-1}{2} \rceil + r)k}$
$n = rk,$ $r = 2h + 1, \ h \geq 2$	$S_4(rk, k, q) = h \left((q^k - 1)^h (q^{rk} - 1) + \frac{(q^k - 1)^{h-1} (q^{rk} - 1)}{q - 1} \right)$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$
$n = rk,$ $r = 2h + 1, \ h \geq 2$	$S_5(rk, k, q) = h q^k (q^k - 1)^{h-1} (q^{rk} - 1) + \frac{q^{rk}-1}{q^k-1}$	$\sim q^{(\lfloor \frac{r-1}{2} \rfloor + r)k}$

Conclusions

Before our work

$$\lim_{k(\text{Or } q) \rightarrow +\infty} \frac{S(2k, k, q)}{J_q(2k, 2k-2, k)} = \frac{1}{2} \quad \text{(Almost)}$$

$n = 2k$ Asymptotically optimal cyclic subspace codes

R. M. Roth, N. Raviv, & I. Tamo. Construction of Sidon spaces with applications to coding. *IEEE Transactions on Information Theory*, 64(6), 4412-4422, **2018**.

$$\lim_{k(\text{Or } q) \rightarrow +\infty} \frac{S(4k, k, q)}{J_q(4k, 2k-2, k)} = \frac{1}{2} \quad \text{(Almost)}$$

$n = 4k$ Asymptotically optimal cyclic subspace codes

S. Yu, L. Ji. Two new constructions of cyclic subspace codes via Sidon spaces. *Designs, Codes and Cryptography*, 92(11), 3799-3811, **2024**.

$$\lim_{k(\text{Or } q) \rightarrow +\infty} \frac{S(3k, k, q)}{J_q(3k, 2k-2, k)} = 1 \quad \text{(Almost)}$$

$n = 3k$ Asymptotically optimal cyclic subspace codes

Zhang, H., Tang, C., & Cao, X. Large optimal cyclic subspace codes. *Discrete Mathematics*, 347(7), 114007, **2024**.

Our contribution

C.C., P. Santonastaso. Asymptotically optimal cyclic subspace codes, arXiv: 2025

$$\lim_{k(\text{Or } q) \rightarrow +\infty} \frac{S(2^e 3^h k, k, q)}{J_q(2^e 3^h k, 2k-2, k)} = \frac{1}{2^e}, e \geq 1, h \geq 0 \quad \text{(Almost)}$$

$n = 2^e 3^h k$
 $e \geq 1, h \geq 0$ Asymptotically optimal cyclic subspace codes

$$\lim_{k(\text{Or } q) \rightarrow +\infty} \frac{S(3^e k, k, q)}{J_q(3^e k, 2k-2, k)} = 1, e \geq 2$$

$n = 3^e k$
 $e \geq 2$ Asymptotically optimal cyclic subspace codes

Conclusions

Before our work

$$\lim_{k(\text{Or } q) \rightarrow +\infty} \frac{S(2k, k, q)}{J_q(2k, 2k-2, k)} = \frac{1}{2} \quad \text{(Almost) Asymptotically optimal cyclic subspace codes}$$

$n = 2k$

R. M. Roth, N. Raviv, & I. Tamo. Construction of Sidon spaces with applications to coding. *IEEE Transactions on Information Theory*, 64(6), 4412-4422, **2018**.

$$\lim_{k(\text{Or } q) \rightarrow +\infty} \frac{S(4k, k, q)}{J_q(4k, 2k-2, k)} = \frac{1}{2} \quad \text{(Almost) Asymptotically optimal cyclic subspace codes}$$

$n = 4k$

S. Yu, L. Ji. Two new constructions of cyclic subspace codes via Sidon spaces. *Designs, Codes and Cryptography*, 92(11), 3799-3811, **2024**.

$$\lim_{k(\text{Or } q) \rightarrow +\infty} \frac{S(3k, k, q)}{J_q(3k, 2k-2, k)} = 1 \quad \text{Asymptotically optimal cyclic subspace codes}$$

$n = 3k$

Zhang, H., Tang, C., & Cao, X. Large optimal cyclic subspace codes. *Discrete Mathematics*, 347(7), 114007, **2024**.

Our contribution

C.C., P. Santonastaso. Asymptotically optimal cyclic subspace codes, arXiv: 2025

$$\lim_{k(\text{Or } q) \rightarrow +\infty} \frac{S(2^e 3^h k, k, q)}{J_q(2^e 3^h k, 2k-2, k)} = \frac{1}{2^e}, e \geq 1, h \geq 0 \quad \text{(Almost) Asymptotically optimal cyclic subspace codes}$$

$n = 2^e 3^h k$
 $e \geq 1, h \geq 0$

$$\lim_{k(\text{Or } q) \rightarrow +\infty} \frac{S(3^e k, k, q)}{J_q(3^e k, 2k-2, k)} = 1, e \geq 2 \quad \text{Asymptotically optimal cyclic subspace codes}$$

$n = 3^e k$
 $e \geq 2$

Thank you for your attention!