

Erdős-Ko-Rado sets on the hyperbolic quadrics
 $Q^+(4n+1, q)$

Leo Storme

Ghent University
Department of Mathematics
Belgium
(joint work with Laure Schelfhout)

Irsee 2025

OUTLINE

1 HYPERBOLIC QUADRICS

2 ERDŐS-KO-RADO PROBLEM

OUTLINE

1 HYPERBOLIC QUADRICS

2 ERDŐS-KO-RADO PROBLEM

HYPERBOLIC QUADRICS

- Hyperbolic quadric $\mathcal{Q}^+(2n+1, q)$:
 $X_0X_1 + X_2X_3 + \cdots + X_{2n}X_{2n+1} = 0$,
- $\mathcal{Q}^+(2n+1, q)$ has points, lines, $\dots$, n -spaces,
- n -spaces on $\mathcal{Q}^+(2n+1, q)$ are called *generators*,
- $\mathcal{Q}^+(2n+1, q)$ has $2(q+1)(q^2+1)\cdots(q^n+1)$ generators.

GENERATORS ON HYPERBOLIC QUADRIC

- Set of generators Ω on $\mathcal{Q}^+(2n+1, q)$ can be partitioned into two equivalence classes Ω_1 and Ω_2 :

$$\Pi_1 \sim \Pi_2 \Leftrightarrow \dim(\Pi_1 \cap \Pi_2) \equiv n \pmod{2}.$$

Hyperbolic quadric $Q^+(3, q)$



Millenáris Park, Budapest

GENERATORS ON HYPERBOLIC QUADRIC

- Set of generators Ω on $\mathcal{Q}^+(2n+1, q)$ can be partitioned into two equivalence classes Ω_1 and Ω_2 :

$$\Pi_1 \sim \Pi_2 \Leftrightarrow \dim(\Pi_1 \cap \Pi_2) \equiv n \pmod{2}.$$

- Ω_1 : **Latin generators**
 Ω_2 : **Greek generators**
- $(q+1)(q^2+1)\cdots(q^n+1)$ Latin and Greek generators.
- For $\mathcal{Q}^+(4n+1, q)$, generators of Ω_1 pairwise intersect.
- For $\mathcal{Q}^+(4n+1, q)$, generators of Ω_2 pairwise intersect.

OUTLINE

1 HYPERBOLIC QUADRICS

2 ERDŐS-KO-RADO PROBLEM



P. Erdős



C. Ko



R. Rado

ERDŐS-KO-RADO PROBLEM

Problem: What are largest sets of k -sets in n -set, pairwise intersecting in at least one element?

THEOREM (ERDŐS-KO-RADO)

If S is set of k -sets in n -set Ω , with $2k \leq n$, pairwise intersecting in at least one element, then $|S| \leq \binom{n-1}{k-1}$. If $2k+1 \leq n$, then equality only holds if S consists of all k -sets through fixed element of Ω .

$n = 2k$: If $n = 2k$, other sets with equality: all k -sets in fixed subset of size $n - 1 = 2k - 1$ of Ω .

q -ANALOG OF ERDŐS-KO-RADO PROBLEM

Problem: What are largest sets of $(k - 1)$ -subspaces in $PG(n - 1, q)$, pairwise intersecting non-trivially?

THEOREM (HSIEH, FRANKL-WILSON, NEWMAN-GODSIL)


Let S be set of $(k - 1)$ -subspaces in $PG(n - 1, q)$, with $2(k - 1) + 1 \leq n - 1$, pairwise intersecting non-trivially.

- If $2(k - 1) + 1 < n - 1$, then largest examples consist of all $(k - 1)$ -spaces through fixed point.*
- If $2(k - 1) + 1 = n - 1$, then largest examples consist of all $(k - 1)$ -spaces through fixed point, or all $(k - 1)$ -spaces in given hyperplane of $PG(n - 1 = 2k - 1, q)$.*

ERDŐS-KO-RADO PROBLEM IN FINITE CLASSICAL POLAR SPACES

Problem:

- What are largest sets of generators in finite classical polar space P , pairwise intersecting non-trivially?
- All generators of P through fixed point (point-pencil = p.-p.).





V. Pepe and F. Vanhove

RESULTS FOR FINITE CLASSICAL POLAR SPACES

Polar space	Maximum size	Classification
$Q^-(2n+1, q)$	$(q^2+1) \cdots (q^n+1)$	p.-p.
$Q(4n, q)$	$(q+1) \cdots (q^{2n-1}+1)$	p.-p.
$Q(4n+2, q), n \geq 2$	$(q+1) \cdots (q^{2n}+1)$	p.-p., Latins $Q^+(4n+1, q)$
$Q(6, q)$	$(q+1)(q^2+1)$	p.-p., Latins $Q^+(5, q)$, base
$Q^+(4n+1, q)$	$(q+1) \cdots (q^{2n}+1)$	all Latins
Latins $Q^+(4n+3, q), n \geq 2$	$(q+1) \cdots (q^{2n}+1)$	p.-p.
Latins $Q^+(7, q)$	$(q+1)(q^2+1)$	p.-p., meeting Greek in plane
$W(4n+1, q), n \geq 2, q$ odd	$(q+1) \cdots (q^{2n}+1)$	p.-p.
$W(4n+1, q), n \geq 2, q$ even	$(q+1) \cdots (q^{2n}+1)$	p.-p., Latins $Q^+(4n+1, q)$
$W(5, q), q$ odd	$(q+1)(q^2+1)$	p.-p., base,
$W(5, q), q$ even	$(q+1)(q^2+1)$	p.-p., base, Latins $Q^+(5, q)$
$W(4n+3, q)$	$(q+1) \cdots (q^{2n+1}+1)$	p.-p.
$H(2n, q^2)$	$(q^3+1)(q^5+1) \cdots (q^{2n-1}+1)$	p.-p.
$H(4n+3, q^2)$	$(q+1)(q^3+1) \cdots (q^{4n+1}+1)$	p.-p.
$H(4n+1, q^2), n \geq 2$	$< \Omega /(q^{2n+1}+1)$	?
$H(5, q^2)$	$q(q^4+q^2+1)+1$	base

LARGEST EKR-SETS ON $Q^+(4n+1, q)$

THEOREM

The largest Erdős-Ko-Rado sets on the hyperbolic quadric $Q^+(4n+1, q)$ are the set of Latin and the set of Greek generators of $Q^+(4n+1, q)$.

This example has size $(q+1)(q^2+1)\cdots(q^{2n}+1) \in O(q^{2n^2+n})$.

Comparison with point-pencil:

Point-pencil on $Q^+(4n+1, q)$ has size

$$2(q+1)(q^2+1)\cdots(q^{2n-1}+1) \in O(q^{2n^2-n}).$$

CLASSIFICATION OF EKR SETS ON $Q^+(5, q)$

THEOREM

There are three distinct maximal EKR-sets on the hyperbolic quadric $Q^+(5, q)$:

- 1 *One class of generators.*
- 2 *Set consisting of one fixed generator Π and all generators of the other class intersecting this fixed generator in at least one point.*
- 3 *Point-pencil: all generators through a fixed point.*

SECOND LARGEST EKR-SET ON $\mathcal{Q}^+(4n+1, q)$


THEOREM (DE BOECK)

The second largest EKR-sets on $\mathcal{Q}^+(4n+1, q)$ are equal to the sets of generators consisting of one fixed generator Π , and the set of generators of the other class, intersecting this fixed generator Π in at least one point.

Class of generators: size $\in O(q^{2n^2+n})$.

Second largest example: size $\in O(q^{2n^2+n-1})$.

SECOND LARGEST EKR-SET ON $Q^+(4n+1, q)$



THIRD LARGEST EKR-SET ON $\mathcal{Q}^+(4n+1, q)$

THEOREM (SCHELFHOUT AND STORME)

The third largest maximal EKR-sets on $\mathcal{Q}^+(4n+1, q)$ are constructed in the following way (up to interchange of Latin and Greek):

- *Consider the $q+1$ Latin generators Π_i , $i = 1, \dots, q+1$, through a fixed $(2n-2)$ -dimensional space Ω .*
- *Consider the set of all Greek generators Π intersecting Ω in at least one point.*

The third largest maximal EKR-sets on $\mathcal{Q}^+(4n+1, q)$ are equal to the sets consisting of these $q+1$ Latin generators Π_i , $i = 1, \dots, q+1$, through a fixed $(2n-2)$ -dimensional space Ω and these Greek generators intersecting in at least one point.

Size $\in O(q^{2n^2+n-2})$.

STEPS IN THE PROOF

- There are at least two generators of each class.
- If all generators of one class pass through a common $(2n - 2)$ -dimensional space, then it the desired example.
- If $|E|$ is at least size of the desired example, then one class of generators contains at least $|E|/2$ generators of S .
- Assume that there are at least $|E|/2$ Greek generators.

STEPS IN THE PROOF

- There are two Greek generators in E that share a $(2n - 2)$ -dimensional space π_{2n-2} .
- But not all Greek generators in E go through π_{2n-2} .
So there is a third Greek generator in E not through π_{2n-2} .
- The number of Latin generators is at most

$$\prod_{i=1}^{2n} (q^i + 1) - q^{2n^2+n} - q^{2n^2+n-1} - q^{2n^2+n-2}.$$

STEPS IN THE PROOF

- The number of Latin generators is at most

$$\prod_{i=1}^{2n} (q^i + 1) - q^{2n^2+n} - q^{2n^2+n-1} - q^{2n^2+n-2}.$$

- No two Latin generators in E share a $(2n - 2)$ -dimensional space, or number of Greek generators is also at most

$$\prod_{i=1}^{2n} (q^i + 1) - q^{2n^2+n} - q^{2n^2+n-1} - q^{2n^2+n-2}.$$

.

Then E is too small.

- So Latin generators in E pairwise intersect in at most $2n - 4$ dimensions.

STEPS IN THE PROOF

- So Latin generators in E pairwise intersect in at most $2n - 4$ dimensions.
- Then E is too small.
- Calculate size of third largest example minus upper bound on size of E leads to

$$q^{2n^2+n-3} - q^{2n^2+n-4} - (q^2 + 1) \prod_{i=4}^{2n} (q^i + 1) + q + 1 > 0.$$

Thank you very much for your attention