

# Representability of uniform $q$ -matroids

Ferdinando Zullo

Università degli Studi della Campania “Luigi Vanvitelli”

(Joint work with G.N. Alfarano, M. Borello, R. Jurrius, A. Neri and O. Polverino)

**Finite Geometries 2025 – Seventh Irsee Conference**

# Matroids vs $q$ -Matroids

## Matroid

$\mathcal{M}$  is a pair  $([n], r)$  where  $r: 2^{[n]} \rightarrow \mathbb{Z}$

$\forall A, B \subseteq [n]$

**(r1) (Boundness)**  $0 \leq r(A) \leq |A|$

**(r2) (Monotonicity)**

$A \subseteq B \Rightarrow r(A) \leq r(B)$

**(r3) (Submodularity)**

$r(A \cup B) + r(A \cap B) \leq r(A) + r(B)$

Rank of  $\mathcal{M}$ :  $r([n])$

## $q$ -Matroid

$\mathcal{M}$  is a pair  $(\mathbb{F}_q^n, \rho)$  where

$\rho: \mathcal{L}(\mathbb{F}_q^n) \rightarrow \mathbb{Z} \quad \forall A, B \in \mathcal{L}(\mathbb{F}_q^n)$

**(r1) (Boundness)**  $0 \leq \rho(A) \leq \dim(A)$

**(r2) (Monotonicity)**

$A \subseteq B \Rightarrow \rho(A) \leq \rho(B)$

**(r3) (Submodularity)**

$\rho(A + B) + \rho(A \cap B) \leq \rho(A) + \rho(B)$

Rank of  $\mathcal{M}$ :  $\rho(\mathbb{F}_q^n)$

Jurrius, R. P. M. J., and G. R. Pellikaan.  
"Defining the  $q$ -analogue of a matroid." The  
Electronic Journal of Combinatorics 25.3

## Matroid

$\mathcal{M}$  is a pair  $([n], r)$  where  $r: 2^{[n]} \rightarrow \mathbb{Z}$   
 $\forall A, B \subseteq [n]$

**(r1) (Boundness)**  $0 \leq r(A) \leq |A|$

**(r2) (Monotonicity)**

$A \subseteq B \Rightarrow r(A) \leq r(B)$

**(r3) (Submodularity)**

$r(A \cup B) + r(A \cap B) \leq r(A) + r(B)$

Rank of  $\mathcal{M}$ :  $r([n])$

## q-Matroid

$\mathcal{M}$  is a pair  $(\mathbb{F}_q^n, \rho)$  where

$\rho: \mathcal{L}(\mathbb{F}_q^n) \rightarrow \mathbb{Z} \quad \forall A, B \in \mathcal{L}(\mathbb{F}_q^n)$

**(r1) (Boundness)**  $0 \leq \rho(A) \leq \dim(A)$

**(r2) (Monotonicity)**

$A \subseteq B \Rightarrow \rho(A) \leq \rho(B)$

**(r3) (Submodularity)**

$\rho(A + B) + \rho(A \cap B) \leq \rho(A) + \rho(B)$

Rank of  $\mathcal{M}$ :  $\rho(\mathbb{F}_q^n)$

Jurrius, R. P. M. J., and G. R. Pellikaan.  
"Defining the q-analogue of a matroid." The  
Electronic Journal of Combinatorics 25.3

## Matroid

$\mathcal{M}$  is a pair  $([n], r)$  where  $r: 2^{[n]} \rightarrow \mathbb{Z}$

$\forall A, B \subseteq [n]$

**(r1) (Boundness)**  $0 \leq r(A) \leq |A|$

**(r2) (Monotonicity)**

$A \subseteq B \Rightarrow r(A) \leq r(B)$

**(r3) (Submodularity)**

$r(A \cup B) + r(A \cap B) \leq r(A) + r(B)$

Rank of  $\mathcal{M}$ :  $r([n])$

## $q$ -Matroid

$\mathcal{M}$  is a pair  $(\mathbb{F}_q^n, \rho)$  where

$\rho: \mathcal{L}(\mathbb{F}_q^n) \rightarrow \mathbb{Z} \quad \forall A, B \in \mathcal{L}(\mathbb{F}_q^n)$

**(r1) (Boundness)**  $0 \leq \rho(A) \leq \dim(A)$

**(r2) (Monotonicity)**

$A \subseteq B \Rightarrow \rho(A) \leq \rho(B)$

**(r3) (Submodularity)**

$\rho(A + B) + \rho(A \cap B) \leq \rho(A) + \rho(B)$

Rank of  $\mathcal{M}$ :  $\rho(\mathbb{F}_q^n)$

Jurrius, R. P. M. J., and G. R. Pellikaan.  
"Defining the  $q$ -analogue of a matroid." The  
Electronic Journal of Combinatorics 25.3

Alfarano, Byrne, Ceria, Gluesing-Luerssen,  
Jany, Jurrius and Pellikaan...

## Matroid

$\mathcal{M}$  is a pair  $([n], r)$  where  $r: 2^{[n]} \rightarrow \mathbb{Z}$

$\forall A, B \subseteq [n]$

**(r1) (Boundness)**  $0 \leq r(A) \leq |A|$

**(r2) (Monotonicity)**

$A \subseteq B \Rightarrow r(A) \leq r(B)$

**(r3) (Submodularity)**

$r(A \cup B) + r(A \cap B) \leq r(A) + r(B)$

## q-Matroid

$\mathcal{M}$  is a pair  $(\mathbb{F}_q^n, \rho)$  where

$\rho: \mathcal{L}(\mathbb{F}_q^n) \rightarrow \mathbb{Z} \quad \forall A, B \in \mathcal{L}(\mathbb{F}_q^n)$

**(r1) (Boundness)**  $0 \leq \rho(A) \leq \dim(A)$

**(r2) (Monotonicity)**

$A \subseteq B \Rightarrow \rho(A) \leq \rho(B)$

**(r3) (Submodularity)**

$\rho(A + B) + \rho(A \cap B) \leq \rho(A) + \rho(B)$

**Example: Uniform matroid**  $U_{k,n}$

$U_{k,n} = ([n], r)$ , where

$r(A) = \min\{|A|, k\}, \quad \forall A \subseteq [n]$

**Example: Uniform q-matroid**  $\mathcal{U}_{k,n}(q)$

$\mathcal{U}_{k,n}(q) = (\mathbb{F}_q^n, \rho)$ , where


$\rho(A) = \min\{\dim(A), k\}, \quad \forall A \in \mathcal{L}(\mathbb{F}_q^n)$

## Example: Matroids from codes

$$G = (G^1 \dots G^n) \in \mathbb{F}_q^{k \times n}$$

$\mathcal{M}[G] = ([n], r)$  **where**

$$r(A) = \text{rk}(G^s : s \in A), \quad \forall A \subseteq [n]$$



$\mathcal{M}[G]$  is a representable matroid


**Hamming metric codes**

## Example: $q$ -Matroids from codes

$$G \in \mathbb{F}_{q^m}^{k \times n}$$

$\mathcal{M}[G] = (\mathcal{L}(\mathbb{F}_q^n), \rho)$  **where**

$\forall A \in \mathcal{L}(\mathbb{F}_q^n)$  **consider**  $\text{colspan}(Y_A) = A$   
 $\rho(A) = \text{rk}(GY_A)$




$\mathcal{M}[G]$  is an  $\mathbb{F}_{q^m}$ -representable  
 $q$ -matroid

**Rank metric codes**

$U_{k,n}$  is representable if  $q$  is large enough


$$G = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ \alpha_1 & \alpha_2 & \cdots & \alpha_n \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_1^{k-1} & \alpha_2^{k-1} & \cdots & \alpha_n^{k-1} \end{pmatrix}$$



Existence of MDS codes/arcs

$\mathcal{U}_{k,n}(q)$  is  $\mathbb{F}_{q^m}$ -representable  $\Leftrightarrow m \geq n$

$$G = \begin{pmatrix} \alpha_1 & \alpha_2 & \cdots & \alpha_n \\ \alpha_1^q & \alpha_2^q & \cdots & \alpha_n^q \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_1^{q^{k-1}} & \alpha_2^{q^{k-1}} & \cdots & \alpha_n^{q^{k-1}} \end{pmatrix}$$



Existence of MRD codes/  
scattered subspaces w.r.t.  
hyperplanes

$U_{k,n}$  is representable if  $q$  is large enough

$$G = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ \alpha_1 & \alpha_2 & \cdots & \alpha_n \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_1^{k-1} & \alpha_2^{k-1} & \cdots & \alpha_n^{k-1} \end{pmatrix}$$

$\mathcal{U}_{k,n}(q)$  is  $\mathbb{F}_{q^m}$ -representable  $\Leftrightarrow m \geq n$

$$G = \begin{pmatrix} \alpha_1 & \alpha_2 & \cdots & \alpha_n \\ \alpha_1^q & \alpha_2^q & \cdots & \alpha_n^q \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_1^{q^{k-1}} & \alpha_2^{q^{k-1}} & \cdots & \alpha_n^{q^{k-1}} \end{pmatrix}$$

There are matroids that are not representable

Vamos, Peter. "On the representation of independence structures." Unpublished manuscript 120 (1968).

There are  $q$ -matroids that are not representable

- \* Byrne, E., Ceria, M., Ionica, S., & Jurrius, R. (2024). Weighted subspace designs from  $q$ -polymatroids. *Journal of Combinatorial Theory, Series A*, 201, 105799.
- \* Gluesing-Luerssen, H., & Jany, B. (2022).  $q$ -Polymatroids and their relation to rank-metric codes. *Journal of Algebraic Combinatorics*, 56(3), 725-753.

# Direct sum of uniform $q$ -matroids

- \*  $\mathcal{M}_i = (\mathbb{F}_q^{n_i}, \rho_i), i = 1, 2$
- \*  $\mathbb{F}_q^n = \mathbb{F}_q^{n_1} \oplus \mathbb{F}_q^{n_2}$
- \*  $\pi_i: \mathbb{F}_q^n \rightarrow \mathbb{F}_q^{n_i}$ , **projections onto the first  $n_1$  and last  $n_2$  coordinates**

- \*  $\mathcal{M}_i = (\mathbb{F}_q^{n_i}, \rho_i), i = 1, 2$
- \*  $\mathbb{F}_q^n = \mathbb{F}_q^{n_1} \oplus \mathbb{F}_q^{n_2}$
- \*  $\pi_i: \mathbb{F}_q^n \rightarrow \mathbb{F}_q^{n_i}$ , **projections onto the first  $n_1$  and last  $n_2$  coordinates**

**Direct sum  $\mathcal{M}_1 \oplus \mathcal{M}_2$  is**

**$(\mathbb{F}_q^n, \rho)$  where**

$$\rho(V) = \dim(V) + \min_{X \leq V} (\rho_1(\pi_1(X)) + \rho_2(\pi_2(X)) - \dim(X))$$

**Ceria, M., & Jurrius, R. (2024). The direct sum of  $q$ -matroids. *Journal of Algebraic Combinatorics*, 59(2), 291-330.**

- \*  $\mathcal{M}_i = (\mathbb{F}_q^{n_i}, \rho_i), i = 1, 2$
- \*  $\mathbb{F}_q^n = \mathbb{F}_q^{n_1} \oplus \mathbb{F}_q^{n_2}$
- \*  $\pi_i: \mathbb{F}_q^n \rightarrow \mathbb{F}_q^{n_i}$ , **projections onto the first  $n_1$  and last  $n_2$  coordinates**

**Direct sum  $\mathcal{M}_1 \oplus \mathcal{M}_2$  is**

**$(\mathbb{F}_q^n, \rho)$  where**

$$\rho(V) = \dim(V) + \min_{X \leq V} (\rho_1(\pi_1(X)) + \rho_2(\pi_2(X)) - \dim(X))$$

Ceria, M., & Jurrius, R. (2024). The direct sum of  $q$ -matroids. *Journal of Algebraic Combinatorics*, 59(2), 291-330.


Gluesing-Luerssen, H., & Jany, B. (2024). Decompositions of  $q$ -matroids using cyclic flats. *SIAM Journal on Discrete Mathematics*, 38(4), 2940-2970.

**Associativity**



**Direct sum of  $t$   $q$ -matroids**

If  $\mathcal{M}_1$  and  $\mathcal{M}_2$  are two  $q$ -matroids  
 $\mathbb{F}_{q^m}$ -representable



$\mathcal{M}_1 \oplus \mathcal{M}_2$  is  $\mathbb{F}_{q^m}$ -representable


$$\mathbb{F}_4 = \{0, 1, \omega, \omega + 1\}$$

$$G = \begin{pmatrix} 1 & \omega & 0 & \omega + 1 \\ 0 & 0 & 1 & \omega \end{pmatrix} \in \mathbb{F}_4^{2 \times 4}$$

$\mathcal{M}_G \oplus \mathcal{M}_G$  is not  $\mathbb{F}_{2^m}$ -representable for every  $m$

Gluesing-Luerssen, H., & Jany, B. (2025). Representability of the direct sum of  $q$ -matroids. *Journal of Algebraic Combinatorics*, 61(4), 51.

If  $\mathcal{M}_1$  and  $\mathcal{M}_2$  are two  $q$ -matroids  
 $\mathbb{F}_{q^m}$ -representable




$\mathcal{M}_1 \oplus \mathcal{M}_2$  is  $\mathbb{F}_{q^m}$ -representable

$$\mathcal{M}_1 = \mathcal{U}_{k_1, n_1}(q) \text{ and } \mathcal{M}_2 = \mathcal{U}_{k_2, n_2}(q)$$

Alfarano, G. N., Jurrius, R., Neri, A., & FZ. Representability of the direct sum of uniform  $q$ -matroids. Accepted for publication in Combinatorial Theory.

If  $\mathcal{M}_1$  and  $\mathcal{M}_2$  are two  $q$ -matroids  
 $\mathbb{F}_{q^m}$ -representable



$\mathcal{M}_1 \oplus \mathcal{M}_2$  is  $\mathbb{F}_{q^m}$ -representable


$$\text{PG}(\mathbb{F}_{q^m}^{k_1+k_2}, \mathbb{F}_{q^m})$$

**Theorem (Alfarano, Jurrius, Neri and FZ)**


$\mathcal{U}_{k_1, n_1}(q) \oplus \mathcal{U}_{k_2, n_2}(q)$  is  $\mathbb{F}_{q^m}$ -representable  $\Leftrightarrow$

There exist  $\mathcal{S}_1 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_1}$  and  $\mathcal{S}_2 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_2}$  of dimension  $n_1$   
 and  $n_2$  (resp.)


$$\text{PG}(\mathbb{F}_{q^m}^{k_1} + \langle 0 \rangle, \mathbb{F}_{q^m})$$



$$\text{PG}(\langle 0 \rangle \oplus \mathbb{F}_{q^m}^{k_2}, \mathbb{F}_{q^m})$$



If  $\mathcal{M}_1$  and  $\mathcal{M}_2$  are two  $q$ -matroids  
 $\mathbb{F}_{q^m}$ -representable



$\mathcal{M}_1 \oplus \mathcal{M}_2$  is  $\mathbb{F}_{q^m}$ -representable


$$\text{PG}(\mathbb{F}_{q^m}^{k_1+k_2}, \mathbb{F}_{q^m})$$

**Theorem (Alfarano, Jurrius, Neri and FZ)**

$\mathcal{U}_{k_1, n_1}(q) \oplus \mathcal{U}_{k_2, n_2}(q)$  is  $\mathbb{F}_{q^m}$ -representable  $\Leftrightarrow$


There exist  $\mathcal{S}_1 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_1}$  and  $\mathcal{S}_2 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_2}$  of dimension  $n_1$   
 and  $n_2$  (resp.)

- $\dim_{\mathbb{F}_q}(\mathcal{S}_1 \cap A) \leq k_1 - 1$  for every hyperplane  $A$  of  $\mathbb{F}_{q^m}^{k_1}$
- $\dim_{\mathbb{F}_q}(\mathcal{S}_2 \cap B) \leq k_2 - 1$  for every hyperplane  $B$  of  $\mathbb{F}_{q^m}^{k_2}$




**Scattered w.r.t. the hyperplanes**

$$\text{PG}(\mathbb{F}_{q^m}^{k_1} + \langle 0 \rangle, \mathbb{F}_{q^m})$$




$$\text{PG}(\langle 0 \rangle \oplus \mathbb{F}_{q^m}^{k_2}, \mathbb{F}_{q^m})$$



Lunardon, G. (2017). MRD-codes and linear sets. Journal of Combinatorial Theory, Series A, 149, 1-20.

Sheekey, J., & Van de Voorde, G. (2020). Rank-metric codes, linear sets, and their duality. Designs, Codes and Cryptography, 88(4), 655-675.

If  $\mathcal{M}_1$  and  $\mathcal{M}_2$  are two  $q$ -matroids  
 $\mathbb{F}_{q^m}$ -representable



$\mathcal{M}_1 \oplus \mathcal{M}_2$  is  $\mathbb{F}_{q^m}$ -representable


$$\text{PG}(\mathbb{F}_{q^m}^{k_1+k_2}, \mathbb{F}_{q^m})$$

**Theorem (Alfarano, Jurrius, Neri and FZ)**


$\mathcal{U}_{k_1, n_1}(q) \oplus \mathcal{U}_{k_2, n_2}(q)$  is  $\mathbb{F}_{q^m}$ -representable  $\Leftrightarrow$

There exist  $\mathcal{S}_1 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_1}$  and  $\mathcal{S}_2 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_2}$  of dimension  $n_1$   
 and  $n_2$  (resp.)

- $\dim_{\mathbb{F}_q}(\mathcal{S}_1 \cap A) \leq k_1 - 1$  for every hyperplane  $A$  of  $\mathbb{F}_{q^m}^{k_1}$
- $\dim_{\mathbb{F}_q}(\mathcal{S}_2 \cap B) \leq k_2 - 1$  for every hyperplane  $B$  of  $\mathbb{F}_{q^m}^{k_2}$
- $\dim_{\mathbb{F}_q}(\mathcal{S}_1 \oplus \mathcal{S}_2 \cap H) \leq k_1 + k_2 - 1$  for every hyperplane  $H$  of  $\mathbb{F}_{q^m}^{k_1+k_2}$  s.t.  $\mathbb{F}_{q^m}^{k_1} \oplus \langle 0 \rangle, \langle 0 \rangle \oplus \mathbb{F}_{q^m}^{k_2} \not\subseteq H$



If  $\mathcal{M}_1$  and  $\mathcal{M}_2$  are two  $q$ -matroids  
 $\mathbb{F}_{q^m}$ -representable



$\mathcal{M}_1 \oplus \mathcal{M}_2$  is  $\mathbb{F}_{q^m}$ -representable


$$\text{PG}(\mathbb{F}_{q^m}^{k_1+k_2}, \mathbb{F}_{q^m})$$

**Theorem (Alfarano, Jurrius, Neri and FZ)**


$\mathcal{U}_{k_1, n_1}(q) \oplus \mathcal{U}_{k_2, n_2}(q)$  is  $\mathbb{F}_{q^m}$ -representable  $\Leftrightarrow$

There exist  $\mathcal{S}_1 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_1}$  and  $\mathcal{S}_2 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_2}$  of dimension  $n_1$   
 and  $n_2$  (resp.)

- $\dim_{\mathbb{F}_q}(\mathcal{S}_1 \cap A) \leq k_1 - 1$  for every hyperplane  $A$  of  $\mathbb{F}_{q^m}^{k_1}$
- $\dim_{\mathbb{F}_q}(\mathcal{S}_2 \cap B) \leq k_2 - 1$  for every hyperplane  $B$  of  $\mathbb{F}_{q^m}^{k_2}$
- $\dim_{\mathbb{F}_q}(\mathcal{S}_1 \oplus \mathcal{S}_2 \cap H) \leq k_1 + k_2 - 1$  for every hyperplane  $H$  of  $\mathbb{F}_{q^m}^{k_1+k_2}$  s.t.  $\mathbb{F}_{q^m}^{k_1} \oplus \langle 0 \rangle, \langle 0 \rangle \oplus \mathbb{F}_{q^m}^{k_2} \not\subseteq H$



If  $\mathcal{M}_1$  and  $\mathcal{M}_2$  are two  $q$ -matroids  
 $\mathbb{F}_{q^m}$ -representable



$\mathcal{M}_1 \oplus \mathcal{M}_2$  is  $\mathbb{F}_{q^m}$ -representable


$$\text{PG}(\mathbb{F}_{q^m}^{k_1+k_2}, \mathbb{F}_{q^m})$$

**Theorem (Alfarano, Jurrius, Neri and FZ)**


$\mathcal{U}_{k_1, n_1}(q) \oplus \mathcal{U}_{k_2, n_2}(q)$  is  $\mathbb{F}_{q^m}$ -representable  $\Leftrightarrow$

There exist  $\mathcal{S}_1 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_1}$  and  $\mathcal{S}_2 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_2}$  of dimension  $n_1$   
 and  $n_2$  (resp.)

- $\dim_{\mathbb{F}_q}(\mathcal{S}_1 \cap A) \leq k_1 - 1$  for every hyperplane  $A$  of  $\mathbb{F}_{q^m}^{k_1}$
- $\dim_{\mathbb{F}_q}(\mathcal{S}_2 \cap B) \leq k_2 - 1$  for every hyperplane  $B$  of  $\mathbb{F}_{q^m}^{k_2}$
- $\dim_{\mathbb{F}_q}(\mathcal{S}_1 \oplus \mathcal{S}_2 \cap H) \leq k_1 + k_2 - 1$  for every hyperplane  $H$  of  $\mathbb{F}_{q^m}^{k_1+k_2}$  s.t.  $\mathbb{F}_{q^m}^{k_1} \oplus \langle 0 \rangle, \langle 0 \rangle \oplus \mathbb{F}_{q^m}^{k_2} \not\subseteq H$



If  $\mathcal{M}_1$  and  $\mathcal{M}_2$  are two  $q$ -matroids  
 $\mathbb{F}_{q^m}$ -representable



$\mathcal{M}_1 \oplus \mathcal{M}_2$  is  $\mathbb{F}_{q^m}$ -representable


$$\text{PG}(\mathbb{F}_{q^m}^{k_1+k_2}, \mathbb{F}_{q^m})$$

**Theorem (Alfarano, Jurrius, Neri and FZ)**


$\mathcal{U}_{k_1, n_1}(q) \oplus \mathcal{U}_{k_2, n_2}(q)$  is  $\mathbb{F}_{q^m}$ -representable  $\Leftrightarrow$

There exist  $\mathcal{S}_1 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_1}$  and  $\mathcal{S}_2 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_2}$  of dimension  $n_1$   
 and  $n_2$  (resp.)

- $\dim_{\mathbb{F}_q}(\mathcal{S}_1 \cap A) \leq k_1 - 1$  for every hyperplane  $A$  of  $\mathbb{F}_{q^m}^{k_1}$
- $\dim_{\mathbb{F}_q}(\mathcal{S}_2 \cap B) \leq k_2 - 1$  for every hyperplane  $B$  of  $\mathbb{F}_{q^m}^{k_2}$
- $\dim_{\mathbb{F}_q}(\mathcal{S}_1 \oplus \mathcal{S}_2 \cap H) \leq k_1 + k_2 - 1$  for every hyperplane  $H$  of  $\mathbb{F}_{q^m}^{k_1+k_2}$  s.t.  $\mathbb{F}_{q^m}^{k_1} \oplus \langle 0 \rangle, \langle 0 \rangle \oplus \mathbb{F}_{q^m}^{k_2} \not\subseteq H$



If  $\mathcal{M}_1$  and  $\mathcal{M}_2$  are two  $q$ -matroids  
 $\mathbb{F}_{q^m}$ -representable



$\mathcal{M}_1 \oplus \mathcal{M}_2$  is  $\mathbb{F}_{q^m}$ -representable

$$\text{PG}(\mathbb{F}_{q^m}^{k_1+k_2}, \mathbb{F}_{q^m})$$


**Theorem (Alfarano, Jurrius, Neri and FZ)**

$\mathcal{U}_{k_1, n_1}(q) \oplus \mathcal{U}_{k_2, n_2}(q)$  is  $\mathbb{F}_{q^m}$ -representable  $\Leftrightarrow$


There exist  $\mathcal{S}_1 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_1}$  and  $\mathcal{S}_2 \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}^{k_2}$  of dimension  $n_1$   
 and  $n_2$  (resp.)

- $\dim_{\mathbb{F}_q}(\mathcal{S}_1 \cap A) \leq k_1 - 1$  for every hyperplane  $A$  of  $\mathbb{F}_{q^m}^{k_1}$
- $\dim_{\mathbb{F}_q}(\mathcal{S}_2 \cap B) \leq k_2 - 1$  for every hyperplane  $B$  of  $\mathbb{F}_{q^m}^{k_2}$
- $\dim_{\mathbb{F}_q}(\mathcal{S}_1 \oplus \mathcal{S}_2 \cap H) \leq k_1 + k_2 - 1$  for every hyperplane  $H$  of  $\mathbb{F}_{q^m}^{k_1+k_2}$  s.t.  $\mathbb{F}_{q^m}^{k_1} \oplus \langle 0 \rangle, \langle 0 \rangle \oplus \mathbb{F}_{q^m}^{k_2} \not\subseteq H$

$\mathcal{S}_1 \oplus \mathcal{S}_2$  is  $(\Lambda_{k_1+k_2-1, k_1, k_2}, k_1 + k_2 - 1)$ -evasive



If  $\mathcal{M}_1$  and  $\mathcal{M}_2$  are two  $q$ -matroids  
 $\mathbb{F}_{q^m}$ -representable



$\mathcal{M}_1 \oplus \mathcal{M}_2$  is  $\mathbb{F}_{q^m}$ -representable

$$\text{PG}(\mathbb{F}_{q^m}^{k_1+k_2}, \mathbb{F}_{q^m})$$

**Theorem (Alfarano, Jurrius, Neri and FZ)**


$\mathcal{U}_{k_1, n_1}(q) \oplus \mathcal{U}_{k_2, n_2}(q)$  is  $\mathbb{F}_{q^m}$ -representable for  $m = m_1 m_2$   
 with  $n_i \leq m_i$  and  $\gcd(m_1, m_2) = 1$

**Construction**


$$\mathcal{S}_1 = \{(x, x^q, \dots, x^{q^{k_1-1}}) : x \in \mathbb{F}_{q^{n_1}}\}$$

$$\mathcal{S}_2 = \{(y, y^q, \dots, y^{q^{k_2-1}}) : y \in \mathbb{F}_{q^{n_2}}\}$$

$\mathcal{S}_1 \oplus \mathcal{S}_2$  is  $(\Lambda_{k_1+k_2-1, k_1, k_2}, k_1 + k_2 - 1)$ -evasive



If  $\mathcal{M}_1$  and  $\mathcal{M}_2$  are two  $q$ -matroids  
 $\mathbb{F}_{q^m}$ -representable



$\mathcal{M}_1 \oplus \mathcal{M}_2$  is  $\mathbb{F}_{q^m}$ -representable

$$\text{PG}(\mathbb{F}_{q^m}^{k_1+k_2}, \mathbb{F}_{q^m})$$

**Theorem (Alfarano, Jurrius, Neri and FZ)**

$\mathcal{U}_{k_1, n_1}(q) \oplus \mathcal{U}_{k_2, n_2}(q)$  is  $\mathbb{F}_{q^m}$ -representable for  $m = m_1 m_2$   
 with  $n_i \leq m_i$  and  $\gcd(m_1, m_2) = 1$


**Construction**

$$\mathcal{S}_1 = \{(x, x^q, \dots, x^{q^{k_1-1}}) : x \in \mathbb{F}_{q^{n_1}}\}$$

$$\mathcal{S}_2 = \{(y, y^q, \dots, y^{q^{k_2-1}}) : y \in \mathbb{F}_{q^{n_2}}\}$$


$\mathcal{S}_1 \oplus \mathcal{S}_2$  is  $(\Lambda_{k_1+k_2-1, k_1, k_2}, k_1 + k_2 - 1)$ -evasive

**Extension to**  $\mathcal{U}_{k_1, n_1}(q) \oplus \dots \oplus \mathcal{U}_{k_t, n_t}(q)$




**Characterization of  $m$  for which  $\mathcal{M}$  is  $\mathbb{F}_{q^m}$ -representable ?**

Characterization of  $m$  for which  $\mathcal{M}$  is  $\mathbb{F}_{q^m}$ -representable



$$\mathcal{U}_{1,n_1}(q) \oplus \mathcal{U}_{1,n_2}(q)$$



**Existence of  $S, T \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}$  s.t.  $L_{S \times T} = \{ \langle (s, t) \rangle_{\mathbb{F}_{q^m}} : s \in S, t \in T \}$  and**


$$w_{L_{S \times T}}(\langle (s, t) \rangle_{\mathbb{F}_{q^m}}) = \dim_{\mathbb{F}_q}((S \times T) \cap \langle (s, t) \rangle_{\mathbb{F}_{q^m}}) \leq 1$$

**if  $st \neq 0$**


Napolitano, V., Polverino, O., Santonastaso, P., & Zullo, F. (2022). Linear sets on the projective line with complementary weights. *Discrete Mathematics*, 345(7), 112890.

**Corrado's talk!**

# Characterization of $m$ for which $\mathcal{M}$ is $\mathbb{F}_{q^m}$ -representable



$$\mathcal{U}_{1,n_1}(q) \oplus \mathcal{U}_{1,n_2}(q)$$

 **Existence of  $S, T \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}$  s.t.  $L_{S \times T} = \{ \langle (s, t) \rangle_{\mathbb{F}_{q^m}} : s \in S, t \in T \}$  and**

$$w_{L_{S \times T}}(\langle (s, t) \rangle_{\mathbb{F}_{q^m}}) = \dim_{\mathbb{F}_q}((S \times T) \cap \langle (s, t) \rangle_{\mathbb{F}_{q^m}}) \leq 1$$

**if  $st \neq 0$**


**$m$  is even and  $m \geq 2 \max\{n_1, n_2\}$**

Napolitano, V., Polverino, O., Santonastaso, P., & Zullo, F. (2022). Linear sets on the projective line with complementary weights. *Discrete Mathematics*, 345(7), 112890.

**Corrado's talk!**

# Characterization of $m$ for which $\mathcal{M}$ is $\mathbb{F}_{q^m}$ -representable ?

$$\mathcal{U}_{1,n_1}(q) \oplus \mathcal{U}_{1,n_2}(q)$$

 **Existence of  $S, T \leq_{\mathbb{F}_q} \mathbb{F}_{q^m}$  s.t.  $L_{S \times T} = \{ \langle (s, t) \rangle_{\mathbb{F}_{q^m}} : s \in S, t \in T \}$  and**

$$w_{L_{S \times T}}(\langle (s, t) \rangle_{\mathbb{F}_{q^m}}) = \dim_{\mathbb{F}_q}((S \times T) \cap \langle (s, t) \rangle_{\mathbb{F}_{q^m}}) \leq 1$$

**if  $st \neq 0$**

$m$  is even and  $m \geq 2 \max\{n_1, n_2\}$

Napolitano, V., Polverino, O., Santonastaso, P., & Zullo, F. (2022). Linear sets on the projective line with complementary weights. *Discrete Mathematics*, 345(7), 112890.

**Corrado's talk!**

**Other conditions**


Neri, A., & Stanojkovski, M. (2024). A proof of the Etzion-Silberstein conjecture for monotone and MDS-constructible Ferrers diagrams. *Journal of Combinatorial Theory, Series A*, 208, 105937.

# Characterization of $m$ for which $\mathcal{M}$ is $\mathbb{F}_{q^m}$ -representable ?

$$f(x) = \sum_{i=0}^{m-1} a_i x^{q^i} \in \mathbb{F}_{q^m}[x]$$

$$\mathcal{U}_{1,n_1}(q) \oplus \mathcal{U}_{1,n_2}(q)$$

$$\mathcal{D}(f) = \{(x, f(x), 0) : x \in \mathbb{F}_{q^m}\} \quad \text{PG}(2, q^m)$$




# Characterization of $m$ for which $\mathcal{M}$ is $\mathbb{F}_{q^m}$ -representable ?

$$\mathcal{U}_{1,n_1}(q) \oplus \mathcal{U}_{1,n_2}(q)$$

$$f(x) = x^{q^{n-1}} + x^{q^{2n-1}} + \mu(x^q - x^{q^{n+1}}) \in \mathbb{F}_{q^{2n}}[x]$$

Smaldore V., Zanella C. And FZ. Regular fat linearised polynomials. Ongoing project.

$$\mathcal{D}(f) = \{(x, f(x), 0) : x \in \mathbb{F}_{q^m}\} \quad \text{PG}(2, q^m)$$



# Characterization of $m$ for which $\mathcal{M}$ is $\mathbb{F}_{q^m}$ -representable ?


$$\mathcal{U}_{1,n_1}(q) \oplus \mathcal{U}_{1,n_2}(q)$$

$$f(x) = x^{q^{n-1}} + x^{q^{2n-1}} + \mu(x^q - x^{q^{n+1}}) \in \mathbb{F}_{q^{2n}}[x]$$

$$\mathcal{D}(f) = \{(x, f(x), 0) : x \in \mathbb{F}_{q^m}\} \quad \text{PG}(2, q^m)$$

Smaldore V., Zanella C. And FZ. Regular fat linearised polynomials. Ongoing project.


Santonastaso, P. (2025). Completely decomposable rank-metric codes. Linear Algebra and its Applications 726, 371-404.



# Delsarte duality and direct sum of uniform $q$ -matroids

## Delsarte duality

$L_U$  is an  $\mathbb{F}_q$ -linear set of rank  $n$  in  $\text{PG}(k-1, q^m)$




$$\Lambda = \text{PG}(k-1, q^m)$$

Lunardon, G., & Polverino, O. (2004). Translation ovoids in orthogonal polar spaces. *Forum Mathematicum* (Vol. 16, pp. 663–669).

## Delsarte duality

$L_U$  is an  $\mathbb{F}_q$ -linear set of rank  $n$  in  $\text{PG}(k-1, q^m)$




$\Lambda = \text{PG}(k-1, q^m)$

$\text{PG}(n-1, q^m)$

## Delsarte duality


$L_U$  is an  $\mathbb{F}_q$ -linear set of rank  $n$  in  $\text{PG}(k-1, q^m)$



Lunardon, G., & Polverino, O. (2004). Translation ovoids in orthogonal polar spaces. *Forum Mathematicum* (Vol. 16, pp. 663-669).

## Delsarte duality


$L_U$  is an  $\mathbb{F}_q$ -linear set of rank  $n$  in  $\text{PG}(k-1, q^m)$



$$L_U = \text{proj}_{\Gamma, \Lambda}(\Sigma)$$

# Delsarte duality

$L_U$  is an  $\mathbb{F}_q$ -linear set of rank  $n$  in  $\text{PG}(k-1, q^m)$




$\tau' \rightarrow$  **polarity of**  $\text{PG}(n-1, q)$   
 $(\perp')$

$\tau \rightarrow$  **polarity of**  $\text{PG}(n-1, q^m)$   
 $(\perp)$

$\text{PG}(n-1, q^m)$

# Delsarte duality

$L_U$  is an  $\mathbb{F}_q$ -linear set of rank  $n$  in  $\text{PG}(k-1, q^m)$




$\tau' \rightarrow$  **polarity of**  $\text{PG}(n-1, q)$   
 $(\perp')$

$\tau \rightarrow$  **polarity of**  $\text{PG}(n-1, q^m)$   
 $(\perp)$

$\text{PG}(n-1, q^m)$

# Delsarte duality

$L_U$  is an  $\mathbb{F}_q$ -linear set of rank  $n$  in  $\text{PG}(k-1, q^m)$




$\tau' \rightarrow$  **polarity of**  $\text{PG}(n-1, q)$   
( $\perp'$ )

$\tau \rightarrow$  **polarity of**  $\text{PG}(n-1, q^m)$   
( $\perp$ )

# Delsarte duality

$L_U$  is an  $\mathbb{F}_q$ -linear set of rank  $n$  in  $\text{PG}(k-1, q^m)$



$\tau' \rightarrow$  **polarity of**  $\text{PG}(n-1, q)$   
 $(\perp')$

$\tau \rightarrow$  **polarity of**  $\text{PG}(n-1, q^m)$   
 $(\perp)$

$U^d$  is called the Delsarte dual of  $U$


$\text{PG}(n-1, q^m)$

# Delsarte duality


Borello M., Polverino O. And FZ. Delsarte duality on subspaces and applications to rank-metric codes and q-matroids. Ongoing project.

$$\text{PG}(\mathbb{F}_{q^m}^{k_1+k_2}, \mathbb{F}_{q^m})$$

$$\text{PG}(\mathbb{F}_{q^m}^{n_1+n_2-k_1-k_2}, \mathbb{F}_{q^m})$$



Delsarte dual




$$n = n_1 + n_2, \quad k = k_1 + k_2$$

# Delsarte duality


Borello M., Polverino O. And FZ. Delsarte duality on subspaces and applications to rank-metric codes and q-matroids. Ongoing project.

$$\text{PG}(\mathbb{F}_{q^m}^{k_1+k_2}, \mathbb{F}_{q^m})$$

$$\text{PG}(\mathbb{F}_{q^m}^{n_1+n_2-k_1-k_2}, \mathbb{F}_{q^m})$$



Delsarte dual



$(\Lambda_{k-1, k_1, k_2}, k-1)$ -evasive

$(\Lambda_{n-k-1, n_1-k_1, n_2-k_2}, n-k-1)$ -evasive


$$n = n_1 + n_2, \quad k = k_1 + k_2$$

# Delsarte duality


Borello M., Polverino O. And FZ. Delsarte duality on subspaces and applications to rank-metric codes and  $q$ -matroids. Ongoing project.

$$\text{PG}(\mathbb{F}_{q^m}^{k_1+k_2}, \mathbb{F}_{q^m})$$

$$\text{PG}(\mathbb{F}_{q^m}^{n_1+n_2-k_1-k_2}, \mathbb{F}_{q^m})$$



Delsarte dual



**Theorem (Borello, Polverino and FZ)**

$\mathcal{U}_{k_1, n_1}(q) \oplus \dots \oplus \mathcal{U}_{k_t, n_t}(q)$  is  $\mathbb{F}_{q^m}$ -representable  $\Leftrightarrow \mathcal{U}_{n_1-k_1, n_1}(q) \oplus \dots \oplus \mathcal{U}_{n_t-k_t, n_t}(q)$  is  $\mathbb{F}_{q^m}$ -representable


# In conclusion...

- \* Representability of the direct sum of uniform  $q$ -matroids**
- \* Open: to find the smallest extension field required for the representability of the direct sum of uniform  $q$ -matroids**

# In conclusion...

- \* Representability of the direct sum of uniform  $q$ -matroids

- \* Open: to find the smallest extension field required for the representability of the direct sum of uniform  $q$ -matroids




More on the existence of  $(\Lambda_{k-1,k_1,k_2}, k-1)$ -evasive

# In conclusion...

- \* Representability of the direct sum of uniform  $q$ -matroids

- \* Open: to find the smallest extension field required for the representability of the direct sum of uniform  $q$ -matroids



More on the existence of  $(\Lambda_{k-1,k_1,k_2}, k-1)$ -evasive

- \* Open: what about the direct sum of other classes of representable  $q$ -matroids?

# OpeRa 2026 - Bordeaux

Open Problems on Rank-Metric Codes

February 23-27, 2026

**Save the date!**



## Invited speakers

- **Xavier Caruso** (Université de Bordeaux - France)
- **Alain Couvreur** (INRIA Saclay - France)
- **Anna-Lena Horlemann** (University of St. Gallen - Switzerland)
- **Relinde Jurrius** (Netherlands Defence Academy - the Netherlands)
- **Alberto Ravagnani** (Eindhoven University of Technology - the Netherlands)
- **John Sheekey** (University College Dublin - Ireland)

## Organizers

- **Gianira Alfarano**
- **Elena Berardini**
- **Martino Borello**
- **Ferdinando Zullo**

**Thanks for your attention!**